

S: 30 April 2001

MCCS-FCD-L (25-30xx)

MEMORANDUM FOR SEE DISTRIBUTION

SUBJECT: Review of Initial Draft, Field Manual (FM) 4-25.11, First Aid

1. Provided for your review and comments on our web site at <http://dcdd.amedd.army.mil/index1.htm> is a copy of the Initial Draft of FM 4-25.11 that will replace FM 21-11 when published. In accordance with TRADOC Regulation 25-36, all draft doctrine posted to web sites for staffing are password controlled. The password to access FM 4-25.11 is "meddoc." This manual provides information on first aid for all service members.

2. Request your review and comments be provided this office no later than 15 April 2001. Request comments be categorized as follows:

a. **Critical Comments.** Critical comments **will** cause nonconcurrence in the draft if the concern is not satisfactorily resolved.

b. **Major comments.** Major comments are significant concerns that **may** result in nonconcurrence in the entire document. This category may be used as a general statement of concern within a subject area, but followed by detailed comments on specific entries in the document.

c. **Substantive comments.** Substantive comments are provided because sections in the document appear to be or are potentially incorrect, incomplete, misleading, or confusing.

MCCS-FCD-L

SUBJECT: Review of Initial Draft, Field Manual (FM) 4-25.11, First Aid

3. Division surgeons are requested to provide the information provided in this memorandum to all medical platoons and medical section leaders assigned to their division.

4. The point of contact for this action is Mr. Gary Simons, telephone: DSN 471-9368, commercial 210-221-9368; FAX DSN 471-9662; e-mail address is gary.simons@cen.amedd.army.mil.

FOR THE COMMANDER:

JOHN E. BALL
Colonel, MS
Director, Combat and
Doctrine Development

DISTRIBUTION:

Headquarters United States Air Force, ATTN: SGWN
110 Luke Avenue, Room 400 Bolling AFB, Washington D.C
20332-7050
United States Navy Warfare Development Command,
Doctrine (N55)
686 Cushing Road (Sims Hall)
Newport, RI 02841-1207
United States Marine Corps Development Command,
ATTN: Logistics Doctrine (C-42)
3300 Russell Road
Quantico, VA 22134-5001
Army National Guard Readiness Center, ATTN: NGB-ARS,
111 South George Mason Drive, Arlington, VA 22204-1382
U.S. Army Forces Command, ATTN: AFMD, Fort McPherson,
GA 30330-6000
I Corps and Fort Lewis, ATTN: Corps Surgeon, Fort Lewis,
WA 98433-5000
III Corps and Fort Hood, ATTN: AFZF-GT-DT Fort Hood, TX
76544-5000

V Corps, ATTN: Corps Surgeon, Unit 29355, APO AE 09014
XVIII ABN Corps and Fort Bragg, ATTN: Corps Surgeon,
Fort Bragg, NC 28307-5000

COMMANDER

1st Cavalry Division, ATTN: Division Surgeon, Fort Hood,
TX 76545-6100
1st Infantry Division (MECH), ATTN: Division Surgeon,
APO AE 09036
1st Armored Division, ATTN: Division Surgeon, APO AE
09252
2d Infantry Division, ATTN: Division Surgeon, APO AP
96224
3d Infantry Division (MECH), ATTN: Division Surgeon,
Fort Stewart, GA 31314
4th Infantry Division (MECH), ATTN: Division Surgeon,
Fort Hood, TX 76545-5200
10th Mountain Division (Light), ATTN: Division Surgeon,
Fort Drum, NY 13602-5000
25th Infantry Division (Light), ATTN: Division Surgeon,
Schofield Barracks, HI 96857-6042
28th Infantry Division, ATTN: Division Surgeon, 14th &
Calder Street, Harrisburg, PA 17013
29th Infantry Division (Light), ATTN: Division Surgeon,
BLDG 2247-57, Fort Belvoir, VA 22060-5057
34th Infantry Division, ATTN: Division Surgeon, 13865
South Robert Trail, Rosemount, MN 55068-3438
35th Infantry Division (MECH), ATTN: Division Surgeon,
2 Sherman Ave, Fort Leavenworth, KS 66027-2346
38th Infantry Division, ATTN: Division Surgeon, 3912 W.
Minnesota Street, Indianapolis, IN 46241-0326
40th Infantry Division, ATTN: Division Surgeon, Bldg 16,
MAS, Los Alimitos, CA 90720
42d Infantry Division, ATTN: Division Surgeon,
137 Glenmore Road, Troy, NY 12180-8398
49th Armored Division, ATTN: Division Surgeon, Camp Narby,
P.O. Box 5218, Austin, TX 77054-2025
75th Division (Exercise), ATTN: G3, 1850 Old Spanish
Trail, Houston, TX 77054-2025
78th Division (Exercise), ATTN: G3, 91 Truman Drive,
Edison, NJ 08817-2487
82d Airborne Division, ATTN: Division Surgeon,
Fort Bragg, NC 28307-5100
85th Division (Exercise) ATTN: G3, 1515 W Central Road,
Arlington Heights, IL, 60005-2475
87th Division(Exercise) ATTN: G3, 1400 Golden Acorn
Drive, Birmingham, AL 35244-1295
91st Division (Exercise), ATTN: G3, Bldg 602, Murray
Circle, Fort Baker, CA 94965-5099
101st Airborne Division (Air Assault) ATTN: Division
Surgeon, Fort Campbell, KY 42223-5000

2d Armored Cavalry Regiment, ATTN: Surgeon, Fort Polk,
LA 71459-5000

MCCS--FCD-L

SUBJECT: Review of Initial Draft, Field Manual (FM) 4-
25.11, First Aid

DISTRIBUTION: (CONT)

3d Armored Cavalry Regiment, ATTN: Surgeon, Fort Carson,
CO 80913-5000

11th Armored Cavalry Regiment, ATTN: Surgeon, Fort Irwin,
CA 92310

18th Medical Command, ATTN: EAMC-OPS, Unit 15281, APO AP
96205-0054

U.S. Army Medical Command, ATTN: MCOP-O, 2050 Worth Road,
Fort Sam Houston, TX 78234-6000

U.S. Army Aviation Center & Fort Rucker, ATTN: ATZQ-TDS-DB,
Fort Rucker, AL 36362-5000

U.S. Army Command and General Staff College, ATTN:
ATZL-SWW-L, Bldg 111, 1 Reynolds Avenue, Fort
Leavenworth, Kansas 66027-1352

U.S. Army Combined Arms Support Command, ATTN: ATCL-CDD,
3901 A Avenue, Fort Lee, VA 23801-6000

U.S. Army John F. Kennedy Special Warfare Center and
School, ATTN: ATSU-DT-PDA, Fort Bragg, NC 28307-5000

U.S. Army Ordnance Center and School, ATTN: ATSL-DI-CSD,
Aberdeen Proving Grounds, MD 21005-5201

U.S. Army Signal Center and Fort Gordon, ATTN: ATZH-DTL,
Fort Gordon, GA 30905-5075

U.S. Army Soldier Support Center, ATTN: ATSG-TS (Building
1000), Fort Jackson, SC 29207-7035

U.S. Army Engineer Center, ATTN: ATZA-TD, Building 3200,
Room 252, Fort Leonard Wood, MO 65437-6650

U.S. Army Intelligence Center and Fort Huachuca, ATTN:
ATZS-TDL-D, Bldg 61730, Fort Huachuca, AZ 85613-7000

16th Cavalry Regiment, U.S. Army Armor Center and School,
ATTN: Doctrine, Fort Knox, KY 40121-5220

COMMANDANT

U.S. Army Chemical School, ATTN: ATSN-CM-DD,
401 Engineer Loop Suite 1029
Fort Leonard Wood, MO 65437-8926

U.S. Army Field Artillery School, ATTN: ATSF-DP,
Fort Sill, OK 73503-5600

U.S. Army Infantry School, ATTN: ATSH-ATL, Fort Benning,
GA 31905-5416

U.S. Army Air Defense Artillery School, ATTN: ATSA-TAC-D,
Fort Bliss, TX 79916-7125

U.S. Army Military Police School, ATTN: Doctrine and
Concepts Division, Maneuver Support Center, Fort Leonard
Wood, MO 65437-8926

U.S. Army Transportation School, ATTN: ATSP-TDLP,
Fort Eustis, VA 23604-5399
U.S. Army School of Aviation Medicine, ATTN: HSHA-AVO,
Fort Rucker, AL 36362-5333
MCCS-FCD-L
SUBJECT: Review of Initial Draft, Field Manual (FM) 4-
25.11, First Aid

DISTRIBUTION: (CONT)
U.S. Army Medical Department Activity, ATTN: NTC Surgeon
(SFC Rodriguez) P.O. Box 105109, Fort Irwin CA,
92310-5109

MCCS--FCD-L

SUBJECT: Review of Initial Draft, Field Manual (FM) 4-25.11, First Aid

DISTRIBUTION: (CONT)

MFR: Routing staffing of revised FM.

Gary Simons/Wtr

Proofread by

C, DLB

C, DLD

PREFACE

This manual meets the first aid training needs of individual service members. Because medical personnel will not always be readily available, the nonmedical service members must rely heavily on their own skills and knowledge of life-sustaining methods to survive on the integrated battlefield. This publication outlines both self-aid and aid to other service members (buddy aid). More importantly, it emphasizes prompt and effective action in sustaining life and preventing or minimizing further suffering and disability. First aid is the emergency care given to the sick, injured, or wounded before being treated by medical personnel. The term *first aid* can be defined as "urgent and immediate lifesaving and other measures which can be performed for casualties by nonmedical personnel when medical personnel are not immediately available." Nonmedical service members have received basic first aid training and should remain skilled in the correct procedures for giving first aid. This manual is directed to *all* service members. The procedures discussed apply to all types of casualties and the measures described are for use by both male and female service members.

The term *combat health support (CHS)* is used by the United States (US) Army to denote its system of health care delivery on the battlefield. The term *health service support (HSS)* is used by the US Marine Corps to denote its battlefield health care delivery system. In this publication the systems will be referred to as CHS/HSS.

This publication implements and/or is in consonance with the following North Atlantic Treaty Organization (NATO) International Standardization Agreements (STANAGs) and American, British, Canadian, and Australian Quadripartite Standardization Agreements (QSTAGs).

Title	STANAG	QSTAG
Medical Training in First Aid, Basic Hygiene and Emergency Care	2122	*535
First Aid Kits and Emergency Medical Care Kits	2126	
Medical First Aid and Hygiene Training in NBC Operations	2358	*535
First Aid Material for Chemical Injuries	2871	

*The title for QSTAG 535 is Medical Training in First Aid, Basic Hygiene and Emergency Care; however it is the companion QSTAG for both STANAGs 2122 and 2358.

When amendment, revision, or cancellation of this publication is proposed which will affect or violate the international agreements concerned, the preparing agency will take appropriate reconciliatory action through international standardization channels. These

FM 21-11
Final Draft

agreements are available on request, using Department of Defense (DD) Form 1425 from the Standardization Documents Order Desk, 700 Robins Avenue, Building 4, Section D, Philadelphia, Pennsylvania 19111-5094.

Unless this publication states otherwise, masculine nouns and pronouns do not refer exclusively to men.

Commercial products (trade names or trademarks) mentioned in this publication are to provide descriptive information and for illustrative purposes only. Their use does not imply endorsement by the Department of Defense (DOD).

The proponent for this publication is the United States (US) Army Medical Department Center and School. Submit comments and recommendations for the improvement of this publication on Department of the Army (DA) Form 2028, Recommended Changes to Publications and Blank Forms or by letter. Request comments and recommendations address the specific page and line numbers and a rationale or justification be provided for each comment. Submits comments and recommendations directly to the **Commander, US Army Medical Department Center and School, ATTN: MCCS-FCD-L, 1400 East Grayson Street, Fort Sam Houston, Texas 78234-6175.**

Field Manual
NO. 21-11

Headquarters
Department of the Army
Washington, DC

FIRST AID

TABLE OF CONTENTS

	Page
PREFACE	v
CHAPTER 1. FUNDAMENTAL CRITERIA FOR FIRST AID	
1-1. General.....	1-1
1-2. Terminology.....	1-1
1-3. Understanding Vital Body Functions for First Aid.....	1-2
1-4. Adverse Conditions.....	1-6
1-5. Basics of First Aid	1-7
1-6. Evaluating a Casualty.....	1-7
1-7. Medical Assistance	1-12
CHAPTER 2. BASIC MEASURES FOR FIRST AID	
2-1. General.....	2-1
2-2. Wound Contamination.....	2-1
Section I. Open the Airway and Restore Breathing.....	2-1
2-3. General.....	2-1
2-4. Breathing Process.....	2-1
2-5. Assessment of and Positioning the Casualty	2-2
2-6. Opening the Airway of an Unconscious or Not Breathing Casualty.....	2-3
2-7. Rescue Breathing	2-7
2-8. Heartbeat	2-13
2-9. Airway Obstructions	2-13
2-10. Opening the Obstructed Airway--Conscious Casualty.....	2-14
2-11. Open an Obstructed Airway--Casualty Lying Down or Unconscious.....	2-19
Section II. Stop the Bleeding and Protect the Wound	2-25
2-12. General.....	2-25
2-13. Clothing	2-25
2-14. Entrance and Exit Wounds	2-25
2-15. Field Dressing	2-26
2-16. Manual Pressure.....	2-30

	Page
2-15.	Field Dressing 2-26
2-16.	Manual Pressure..... 2-30
2-17.	Pressure Dressing..... 2-31
2-18.	Tourniquet..... 2-34
Section III.	Check for Shock and Administer First Aid Measures 2-39
2-19.	General..... 2-39
2-20.	Causes and Effects 2-39
2-21.	Signs and Symptoms of Shock 2-29
2-22.	First Aid Measures for Shock 2-40
 CHAPTER 3. FIRST AID FOR SPECIFIC INJURIES	
3-1.	General..... 3-1
3-2.	Head, Neck, and Facial Injuries 3-1
3-3.	General First Aid Measures 3-2
3-4.	Dressings and Bandages..... 3-4
3-5.	Chest Wounds 3-21
3-6.	Chest Wound(s) Procedure 3-21
3-7.	Abdominal Wounds 3-26
3-8.	Abdominal Wound Procedure 3-26
3-9.	Burn Injuries 3-31
3-10.	First Aid for Burns 3-31
3-11.	Shoulder Bandage 3-35
3-12.	Elbow Bandage 3-37
3-13.	Hand Bandage 3-37
3-14.	Leg (Upper and Lower) Bandage..... 3-39
3-15.	Knee Bandage 3-40
3-16.	Foot Bandage 3-41
 CHAPTER 4. FIRST AID FOR FRACTURES	
4-1.	General..... 4-1
4-2.	Kinds of Fractures..... 4-1
4-3.	Signs and Symptoms of Fractures..... 4-2
4-4.	Purposes of Immobilizing Fractures 4-2
4-5.	Splints, Padding, Bandages, Slings, and Swathes..... 4-2
4-6.	Procedures for Splinting Suspected Fractures 4-3
4-7.	Upper Extremity Fractures..... 4-10
4-8.	Lower Extremity Fractures 4-14
4-9.	Jaw, Collarbone, and Shoulder Fractures 4-18
4-10.	Spinal Column Fractures 4-20
4-11.	Neck Fractures 4-23

Page

CHAPTER 5.	FIRST AID FOR CLIMATIC INJURIES	
5-1.	General.....	5-1
5-2.	Heat Injuries.....	5-1
5-3.	Cold Injuries.....	5-4
CHAPTER 6.	FIRST AID FOR BITES AND STINGS	
6-1.	General.....	6-1
6-2.	Types of Snakes	6-1
6-3.	Snakebites	6-6
6-4.	Human and Other Animal Bites.....	6-9
6-5.	Marine (Sea) Animals	6-9
6-6.	Insect (Arthropod) Bites/Stings	6-10
6-7.	Table	6-14
CHAPTER 7.	FIRST AID IN A NUCLEAR, BIOLOGICAL, AND CHEMICAL ENVIRONMENT	
7-1.	General.....	7-1
7-2.	First Aid Materials	7-1
7-3.	Classification of Chemical and Biological Agents	7-2
7-4.	Conditions for Masking Without Order or Alarm	7-2
7-5.	First Aid for a Chemical Attack.....	7-3
7-6.	Background Information on Nerve Agents	7-4
7-7.	Signs and Symptoms of Nerve Agent Poisoning	7-6
7-8.	First Aid for Nerve Agent Poisoning	7-6
7-9.	Blister Agents.....	7-18
7-10.	Choking Agents (Lung-Damaging Agents)	7-19
7-11.	Blood Agents	7-20
7-12.	Incapacitating Agents.....	7-21
7-13.	Incendiaries	7-22
7-14.	Biological Agents and First Aid	7-23
7-15.	Toxins	7-23
7-16.	Nuclear Detonation	7-24
CHAPTER 8.	FIRST AID FOR PSYCHOLOGICAL REACTIONS-- INTRODUCTION	
8-1.	General.....	8-1
8-2.	Importance of Psychological First Aid	8-1
8-3.	Situations Requiring Psychological First Aid.....	8-1
8-4.	Interrelationship of Psychological and Physical First Aid	8-2

	Page
8-5. Goals of Psychological First Aid	8-2
8-6. Respect for Others' Feelings	8-2
8-7. Emotional and Physical Disability.....	8-3
8-8. Battle Fatigue and Other Combat Stress Reactions	8-4
8-9. Reactions to Stress	8-4
8-10. Severe Stress or Battle Fatigue	8-5
8-11. Application of Psychological First Aid.....	8-6
8-12. Reactions and Limitations.....	8-8
8-13. Tables.....	8-8
 APPENDIX A. FIRST AID CASE AND KITS, DRESSINGS, AND BANDAGES	
A-1. First Aid Case with Field Dressings and Bandages	A-1
A-2. General Purpose First Aid Kits	A-2
A-3. Dressings.....	A-2
A-4. Standard Bandages.....	A-2
A-5. Triangular and Cravat (Swathe) Bandages	A-2
 APPENDIX B. RESCUE AND TRANSPORTATION PROCEDURES	
B-1. General.....	B-1
B-2. Principles of Rescue Operations	B-1
B-3. Considerations	B-1
B-4. Plan of Action	B-2
B-5. Proper Handling of Casualties	B-3
B-6. Positioning the Casualty	B-4
B-7. Medical Evacuation and the Transportation of Casualties.....	B-5
B-8. Manual Carries.....	B-6
B-9. Improvised Litters.....	B-32
 APPENDIX C. DIGITAL PRESSURE	C-1
 GLOSSARY	Glossary-1
 REFERENCES	References-1
 INDEX	Index-1

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45

CHAPTER 1

FUNDAMENTAL CRITERIA FOR FIRST AID

1-1. General

Service members may have to depend upon their first aid knowledge and skills to save themselves (self-aid) or other service members (buddy aid). They may be able to save a life, prevent permanent disability, or reduce long periods of hospitalization by knowing **WHAT** to do, **WHAT NOT** to do, and **WHEN** to seek medical assistance.

1-2. Terminology

To enhance the understanding of the material contained in this publication, the following terms are used--

- *Combat lifesaver.* This is a United States (US) Army program governed by Department of the Army (DA) Pamphlet 350-21, *The Combat Lifesaver*. The combat lifesaver is a member of a nonmedical unit selected by the unit commander for additional training beyond basic first aid procedures (referred to as *enhanced first aid*). A minimum of one individual per squad, crew, team, or equivalent-sized unit should be trained. The primary duty of this individual does not change. The additional duty of combat lifesaver is to provide enhanced first aid for injuries based on his training before the combat medic arrives. The combat lifesaver's training is normally provided by medical personnel assigned, attached, or in direct support (DS) of the unit. The senior medical person designated by the commander manages the training program.

- *Combat medic (US Army) or Hospital Corpsman (US Marine Corps).* A medical specialist trained in emergency medical treatment (EMT) procedures and assigned or attached in support of a combat or combat support unit or marine forces.

- *Casualty evacuation.* Casualty evacuation (CASEVAC) is a term used by nonmedical units to refer to the movement of casualties aboard nonmedical vehicles or aircraft. See also the term *transported* below. Refer to FM 8-10-6 for additional information.

CAUTION

Casualties transported in this manner do not receive en route medical care.

- *Enhanced first aid (US Army).* Enhanced first aid is administered by the combat lifesaver. It includes measures, which require an additional level training above self-aid and buddy aid, such as the initiation of IV fluids.

1 • *Evacuation (or medical evacuation).* Medical evacuation is the timely, efficient
2 movement of the wounded, injured, or ill service members from the battlefield and other
3 locations to medical treatment facilities (MTFs). En route medical care is provided by medical
4 personnel during the evacuation. Once the casualty has entered the medical stream (combat
5 medic, hospital corpsman, evacuation crew, or MTF), the role of first aid in the care of the
6 casualty ceases and the casualty becomes the responsibility of the CHS/HSS chain. Once he has
7 entered the CHS/HSS chain he is referred to as a *patient*.

8
9 • *First aid measures.* Urgent and immediate lifesaving and other measures which
10 can be performed for casualties (or performed by the victim himself) by nonmedical personnel
11 when medical personnel are not immediately available.

12
13 • *Medical treatment.* Medical treatment is the care and management of wounded,
14 injured, or ill service members by medically trained (military occupational specialty [MOS]
15 trained, hospital corpsman [HM], and area of concentration [AOC]) personnel). It includes
16 EMT, advanced trauma management (ATM), resuscitative, surgical, definitive, and rehabilitative
17 care provided on the battlefield and in MTFs.

18
19 • *Medical treatment facility.* Any facility established for the purpose of providing
20 medical treatment. This includes battalion aid stations, division clearing stations, dispensaries,
21 clinics, and hospitals.

22
23 • *Self-aid/buddy aid.* Each individual service member is trained to be proficient in
24 a variety of specific first aid procedures. This training enables the service member or a buddy to
25 apply immediate first aid measures to alleviate a life-threatening situation.

26
27 • *Transported.* A casualty is moved to an MTF in a nonmedical vehicle without en
28 route care provided by a medically-trained service member (such as a combat medic or hospital
29 corpsman). First aid measures should be continually performed while the casualty is being
30 transported. If the casualty is acquired by a dedicated medical vehicle with a medically-trained
31 crew, the role of first aid ceases and the casualty becomes the responsibility of the CHS/HSS
32 chain, and is then referred to as a *patient*. This method of transporting a casualty is also referred
33 to as *CASEVAC*.

34 35 **1-3. Understanding Vital Body Functions for First Aid**

36
37 In order for the service member to learn to perform first aid procedures, he must have a basic
38 understanding of what are the vital body functions and what the result will be if they are damaged
39 or not functioning.

40
41 *a. Respiration.* Respiration occurs when a person inhales (oxygen is taken into the
42 body) and then exhales (carbon dioxide [CO₂] is expelled from the body). Respiration involves
43 the--
44

1 • *Airway.* The airway consists of the nose, mouth, throat, voice box, and
2 windpipe. It is the canal through which air passes to and from the lungs.

3
4 • *Lungs.* The lungs are two elastic organs made up of thousands of tiny air
5 spaces and covered by an airtight membrane. The *bronchial tree* is a part of the lungs.

6
7 • *Chest cage.* The chest cage is formed by the muscle-connected ribs, which
8 join the spine in back, and the breastbone in front. The top part of the chest cage is closed by the
9 structure of the neck, and the bottom part is separated from the abdominal cavity by a large
10 dome-shaped muscle called the *diaphragm* (Figure 1-1). The diaphragm and rib muscles, which
11 are under the control of the respiratory center in the brain, automatically *contract* and *relax*.
12 *Contraction* increases and *relaxation* decreases the size of the chest cage. When the chest cage
13 increases and then decreases, the air pressure in the lungs is first less and then more than the
14 atmospheric pressure, thus causing the air to rush into and out of the lungs to equalize the
15 pressure. This cycle of inhaling and exhaling is repeated about 12 to 18 times per minute.
16
17

18 *Figure 1-1. Airway, lungs, and chest cage.*

19
20
21 *b. Blood Circulation.* The heart and the blood vessels (arteries, veins, and
22 capillaries) circulate blood through the body tissues. The heart is divided into two separate
23 halves, each acting as a pump. The left side pumps oxygenated blood (bright red) through the
24 arteries into the capillaries; nutrients and oxygen pass from the blood through the walls of the
25 capillaries into the cells. At the same time waste products and CO₂ enter the capillaries. From
26 the capillaries the oxygen poor blood is carried through the veins to the right side of the heart and
27 then into the lungs where it expels the CO₂ and picks up oxygen. Blood in the veins is dark red
28 because of its low oxygen content. Blood does not flow through the veins in spurts as it does
29 through the arteries. The entire system of the heart, blood vessels, and lymphatics is called the
30 *vascular system*.

31
32 (1) *Heartbeat.* The heart functions as a pump to circulate the blood
33 continuously through the blood vessels to all parts of the body. It contracts, forcing the blood
34 from its chambers; then it relaxes, permitting its chambers to refill with blood. The rhythmical
35 cycle of contraction and relaxation is called the *heartbeat*. The normal heartbeat is from 60 to 80
36 beats per minute.

37
38 (2) *Pulse.* The heartbeat causes a rhythmical expansion and contraction of the
39 arteries as it forces blood through them. This cycle of expansion and contraction can be felt
40 (monitored) at various points in the body and is called the *pulse*. The common points for
41 checking the pulse are at the--

- 42
43 • Side of the neck (*carotid*).
44
45 • Groin (*femoral*).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44

- Wrist (*radial*).
- Ankle (*posterior tibial*).

(a) *Carotid pulse.* To check the carotid pulse, feel for a pulse on the side of the casualty's neck closest to you. This is done by placing the tips of your first two fingers beside his Adam's apple (Figure 1-2).

Figure 1-2. Carotid pulse.

(b) *Femoral pulse.* To check the femoral pulse, press the tips of two fingers into the middle of the groin (Figure 1-3).

Figure 1-3. Femoral pulse.

(c) *Radial pulse.* To check the radial pulse, place your first two fingers on the thumb side of the casualty's wrist (Figure 1-4).

Figure 1-4. Radial pulse.

(d) *Posterior tibial pulse.* To check the posterior tibial pulse, place your first two fingers on the inside of the ankle (Figure 1-5).

Figure 1-5. Posterior tibial pulse.

NOTE

DO NOT use your thumb to check a casualty's pulse because you may confuse the beat of your pulse with that of the casualty.

1-4. Adverse Conditions

a. *Lack of Oxygen.* Human life cannot exist without a continuous intake of oxygen. Lack of oxygen rapidly leads to death. First aid involves knowing how to open the airway and restore breathing.

1 **b. *Bleeding.*** Human life cannot continue without an adequate volume of blood
2 circulating through the body to carry oxygen to the tissues. An important first aid measure is to
3 stop the bleeding to prevent the loss of blood.
4

5 **c. *Shock.*** Shock means there is an inadequate blood flow to the vital tissues and
6 organs. Shock that remains uncorrected may result in death even though the injury or condition
7 causing the shock would not otherwise be fatal. First aid includes preventing shock, since the
8 casualty's chances of survival are much greater if he does not develop shock. Refer to paragraphs
9 2-19 through 2-22 for a further discussion of shock.
10

11 **d. *Infection.*** Recovery from a severe injury or a wound depends largely upon how
12 well the injury or wound was initially protected. Infections result from the multiplication and
13 growth (spread) of harmful microscopic organisms (sometimes referred to as germs). These
14 harmful microscopic organisms are in the air, water, and soil and on the skin and clothing. Some
15 of these organisms will immediately invade (contaminate) a break in the skin or an open wound.
16 The objective is to keep wounds clean and free of these organisms. A good working knowledge
17 of basic first aid measures also includes knowing how to dress a wound to avoid infection or
18 additional contamination.
19

20 **1-5. Basics of First Aid**

21
22 Anything service members can do to keep others in good fighting condition is part of the primary
23 mission to fight or to support the human component of the Army's weapons systems. Most
24 injured or ill service members are able to return to their units to fight or support primarily
25 because they are given appropriate and timely first aid followed by the best medical care
26 possible. Therefore, all service members must remember the basics.
27

- 28 • Check for **BREATHING**: Lack of oxygen intake (through a compromised airway
29 or inadequate breathing) can lead to brain damage or death in very few minutes.
30
- 31 • Check for **BLEEDING**: Life cannot continue without an adequate volume of
32 blood to carry oxygen to tissues.
33
- 34 • Check for **SHOCK**: Unless shock is prevented, first aid performed. and medical
35 treatment provided, death may result even though the injury would not otherwise be fatal.
36

37 **1-6. Evaluating a Casualty**

38
39 **a.** The time may come when you must instantly apply your knowledge of first aid
40 measures. This could occur during combat operations, in training situations, or while in a
41 nonduty status. Any service member observing an unconscious and/or ill, injured, or wounded
42 person must carefully and skillfully evaluate him to determine the first aid measures required to
43 prevent further injury or death. He should seek help from medical personnel as soon as possible,
44 but should not interrupt his evaluation of the casualty or fail to administer first aid measures. A
45 second service member may be sent to find medical help. One of the principles for assisting a

1 casualty is that you (the initial rescuer) must continue the evaluation and first aid measures, as
2 the tactical situation permits, until you are relieved by another individual. If, during any part of
3 the evaluation, the casualty exhibits the conditions (such as shock) for which the service member
4 is checking, the service member must stop the evaluation and immediately administer first aid.
5 In a chemical environment, the service member should not evaluate the casualty until both the
6 individual and the casualty has been masked. If it is suspected that a nerve agent was used,
7 administer the casualty's own nerve agent antidote autoinjector. ***Do not use your own nerve***
8 ***agent antidote autoinjector on the casualty.*** After providing first aid, the service member must
9 proceed with the evaluation and continue to monitor the casualty for further medical
10 complications until relieved by medical personnel.

11
12 **NOTE**

13 Remember, when evaluating and/or administering first aid to a casualty,
14 you should seek medical aid as soon as possible. **DO NOT** stop first aid
15 measures, but if the situation allows, send another service member to find medical
16 aid.

17
18 **WARNING**

19 Again, remember, if there are any signs of a chemical or biological agent
20 attack or poisoning, you should immediately mask and then mask the casualty.
21 If it is suspected to be nerve agent poisoning, administer the antidote using the
22 casualty's autoinjector.

23
24 *b.* To evaluate a casualty, perform the following steps:

25
26 (1) *Check the casualty for responsiveness.* This is done by gently shaking or
27 tapping him while calmly asking, "Are you okay?" Watch for a response. If the casualty does
28 not respond, go to step (2). If the casualty responds, continue with the evaluation.

29
30 (a) If the casualty is conscious, ask him where he feels different than
31 usual or where it hurts. Ask him to identify the location of pain if he can, or to identify the area
32 in which there is no feeling.

33
34 (b) If the casualty is conscious but is choking and cannot talk, stop the
35 evaluation and begin first aid measures. Refer to paragraphs 2-10 and 2-11 for specific
36 information on opening the airway.

37
38 **WARNING**

39 If a broken neck or back is suspected, do not move the casualty unless
40 his life is in immediate danger (such as close to a burning vehicle).
41 Movement may cause permanent paralysis or death.

42
43
44
45 (2) *Check for breathing.* (Refer to paragraph 2-6b[3] for this procedure).

- 1
2 (a) If the casualty is breathing, proceed to step (4).
3
4 (b) If the casualty is not breathing, stop the evaluation and begin first
5 aid measures to attempt to ventilate the casualty. If an airway obstruction is apparent, clear the
6 airway obstruction, then ventilate.
7
8 (c) After successfully ventilating the casualty, proceed to step (3).
9
10 (3) *Check for pulse.* (Refer to paragraph 2-7 for specific methods). If a pulse
11 is present and the casualty is breathing, proceed to step (4).
12
13 (a) If a pulse is present, but the casualty is still not breathing, start
14 rescue breathing.
15
16 (b) If pulse is not present seek medical personnel for help.
17
18 (4) *Check for bleeding.* Look for spurts of blood or blood-soaked clothes.
19 Also check for *both* entry and exit wounds. *If the casualty is bleeding from an open wound,* stop
20 the evaluation and begin first aid procedures as follows for a--
21
22 (a) Wound of the arm or leg (refer to paragraphs 2-15 through 2-17 for
23 information on putting on a field or pressure dressing).
24
25 (b) Partial or complete amputation (refer to paragraph 2-18 for
26 information on putting on a tourniquet).
27
28 (c) Open head wound (refer to paragraph 3-4a for information on
29 applying a dressing to an open head wound).
30
31 (d) Open chest wound (refer to paragraph 3-6 for information on
32 applying a dressing to an open chest wound).
33
34 (e) Open abdominal wound (refer to paragraph 3-8 for information on
35 applying a dressing to an open abdominal wound).
36
37

38 **WARNING**

39 In a chemically contaminated area, do not expose the wounds.
40
41

- 42 (5) *Check for shock.* (Refer to paragraph 2-22 for first aid measures for
43 shock). If the signs and symptoms of shock are present, stop the evaluation and begin first aid
44 measures immediately. The following are the nine signs and symptoms of shock.
45

- 1 (a) Sweaty but cool skin (clammy skin).
2
3 (b) Paleness of skin. (In dark-skinned service members look for a
4 grayish cast to the skin.)
5
6 (c) Restlessness or nervousness.
7
8 (d) Thirst.
9
10 (e) Loss of blood (bleeding).
11
12 (f) Confusion (does not seem aware of surroundings).
13
14 (g) Faster than normal breathing rate.
15
16 (h) Blotchy or bluish skin, especially around the mouth.
17
18 (i) Nausea or vomiting.
19

20 **WARNING**

21 Leg fractures must be splinted before elevating the legs as a first
22 aid measure for shock.

23
24 (6) *Check for fractures.*

- 25
26 (a) Check for the following signs and symptoms of a back or neck
27 injury and perform first aid procedures as necessary.
28
29 • Pain or tenderness of the neck or back area.
30
31 • Cuts or bruises on the neck or back area.
32
33 • Inability of a casualty to move (paralysis or numbness).
34
35 • Ask about ability to move (paralysis).
36
37 • Touch the casualty's arms and legs and ask whether
38 he can feel your hand (numbness).
39
40 • Unusual body or limb position.
41
42 (b) Immobilize any casualty suspected of having a neck or back injury
43 by doing the following:
44
45 • Tell the casualty not to move.

1
2 • If a back injury is suspected, place padding (rolled or folded
3 to conform to the shape of the arch) under the natural arch of the casualty's back. (For example,
4 a blanket may be used as padding.)

5
6 • If a neck injury is suspected, place a roll of cloth under the
7 casualty's neck and put weighted boots (filled with dirt or sand) or rocks on both sides of his
8 head.

9
10 (c) Check the casualty's arms and legs for open or closed fractures.

11
12 • Check for *open* fractures by looking for--

13
14 • Bleeding.

15
16 • Bones sticking through the skin.

17
18 • Check for *closed* fractures by looking for--

19
20 • Swelling.

21
22 • Discoloration.

23
24 • Deformity.

25
26 • Unusual body position.

27
28 (d) Stop the evaluation and begin first aid measures if a fracture to an
29 arm or leg is suspected. Refer to Chapter 4 for information on splinting a suspected fracture.

30
31 (e) Check for signs/symptoms of fractures of other body areas (for
32 example, shoulder or hip) and provide first aid as necessary.

33
34 (7) *Check for burns.* Look carefully for reddened, blistered, or charred skin;
35 also check for singed clothing. If burns are found, stop the evaluation and begin first aid
36 procedures. Refer to paragraph 3-10 for information on giving first aid for burns.

37
38 (8) *Check for possible head injury.*

39
40 (a) Look for the following signs and symptoms:

41
42 • Unequal pupils.

43
44 • Fluid from the ear(s), nose, mouth, or injury site.

45

- 1 • Slurred speech.
- 2
- 3 • Confusion.
- 4
- 5 • Sleepiness.
- 6
- 7 • Loss of memory or consciousness.
- 8
- 9 • Staggering in walking.
- 10
- 11 • Headache.
- 12
- 13 • Dizziness.
- 14
- 15 • Nausea or vomiting.
- 16
- 17 • Paralysis.
- 18
- 19 • Convulsions or twitches.
- 20

21 (b) If a head injury is suspected, continue to watch for signs which
22 would require performance of rescue breathing, first aid measures for shock, or control of
23 bleeding; seek medical aid. Refer to paragraph 3-2 for information on first aid measures for head
24 injuries.

25

26 **1-7. Medical Assistance**

27

28 When a nonmedical service member comes upon an unconscious or injured service member, he
29 must accurately evaluate the casualty to determine the first aid measures needed to prevent
30 further injury or death. He should seek medical assistance as soon as possible, but he should not
31 interrupt the performance of first aid measures. To interrupt the first aid measures may cause
32 more harm than good to the casualty. A second person may be sent to find medical help. If,
33 during any part of the evaluation, the casualty exhibits the conditions (such as shock) for which
34 the service member is checking, the service member must stop the evaluation and immediately
35 administer first aid. Remember that in a chemical environment, the service member should not
36 evaluate the casualty until the casualty has been masked. After performing first aid, the service
37 member must proceed with the evaluation and continue to monitor the casualty for development
38 of conditions which may require the performance of necessary basic life saving measures, such as
39 clearing the airway, rescue breathing, preventing shock, and controlling bleeding. He should
40 continue to monitor the casualty until relieved by medical personnel.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54

CHAPTER 2

BASIC MEASURES FOR FIRST AID

2-1. General

Several conditions that require immediate attention are an inadequate airway, lack of breathing, and excessive loss of blood (circulation). A casualty without a clear airway or who is not breathing may die from lack of oxygen. Excessive loss of blood may lead to shock, and shock can lead to death; therefore, you must act immediately to control the loss of blood.

NOTE

It is also important that you attend to any airway, breathing, or circulation problems **IMMEDIATELY** because these problems, if left unattended, may become life threatening.

2-2. Wound Contamination

All wounds are considered to be contaminated, since infection-producing organisms (germs) are always present on the skin and clothing, and in the soil, water, and air. Any missile or instrument (such as a bullet, shrapnel, knife, or bayonet) causing a wound pushes or carries the germs into that wound. Infection results as these organisms multiply. That a wound is contaminated does not lessen the importance of protecting it from further contamination. You must dress and bandage a wound as soon as possible to prevent further contamination.

2-3. Heartbeat

If a casualty's heart stops beating, you must immediately seek medical personnel for help. **SECONDS COUNT.** Stoppage of the heart is soon followed by cessation of respiration unless it has occurred first. Be calm. Think and act. When a casualty's heart has stopped, there is no pulse at all; the person is unconscious and limp, and the pupils of his eyes are wide open. When evaluating a casualty or when performing the preliminary steps of rescue breathing, feel for a pulse. If you **DO NOT** detect a pulse, immediately seek medical personnel.

Section I. OPEN THE AIRWAY AND RESTORE BREATHING

2-4. General

In order for you to perform first aid, you must understand the breathing process and the consequences if it is interrupted or not functioning. This section discusses the breathing process and what to do if a casualty is having difficulties breathing.

2-5. Breathing Process

All living things must have oxygen to live. Through the breathing process, the lungs draw oxygen from the air and put it into the blood. The heart (which has four chambers) pumps the blood through the body to be used by the cells which require a constant supply of oxygen. Some cells are more dependent on a constant supply of oxygen than others. For example, cells of the brain may die within 4 to 6 minutes without oxygen. Once these cells die, they are lost forever since they do not regenerate. This could result in permanent brain damage, paralysis, or death.

1
2 **2-6. Assessment of and Positioning the Casualty**
3

4 a. **CHECK** for responsiveness (Figure 2-1A)--establish whether the casualty is
5 conscious by gently shaking him and asking, "Are you O.K.?"
6

7 b. **CALL** for help (Figure 2-1B).
8

9 c. **POSITION** the unconscious casualty so that he is lying on his back and on a firm
10 surface (Figure 2-1C).
11

12 **WARNING**

13 If the casualty is lying on his chest (prone position), *cautiously*
14 *roll the casualty as a unit so that his body does not twist* (which
15 may further complicate a neck, back, or spinal injury).
16

17
18 *Figure 2-1. Assessment.*
19
20

21 (1) Straighten the casualty's legs. Take the casualty's arm that is nearest to
22 you and move it so that it is straight and above his head. Repeat the procedure for the other arm.
23

24 (2) Kneel beside the casualty with your knees near his shoulders (leave space
25 to roll his body) (Figure 2-1B). Place one hand behind his head and neck for support. With your
26 other hand, grasp the casualty under his far arm (Figure 2-1C).
27

28 (3) Roll the casualty towards you using a steady, even pull. His head and neck
29 should stay in line with his back.
30

31 (4) Return the casualty's arms to his side. Straighten his legs. Reposition
32 yourself so that you are now kneeling at the level of the casualty's shoulders. However, if a neck
33 injury is suspected and the *jaw-thrust* technique (paragraph 2-6) will be used, kneel at the
34 casualty's head, looking towards his feet.
35

36
37 **2-7. Opening the Airway of an Unconscious or Not Breathing Casualty**
38

39 The tongue is the single most common cause of an airway obstruction (Figure 2-2). In most
40 cases, the airway can be cleared by simply using the *head-tilt/chin-lift* technique. This action
41 pulls the tongue away from the air passage in the throat (Figure 2-3).
42

43 **NOTE**

44 If foreign material or vomitus is visible in the mouth, it should be
45 removed, but do not spend an excessive amount of time doing so.
46

47
48 *Figure 2-2. Airway blocked by tongue.*
49

50
51 *Figure 2-3. Airway opened by extending neck.*
52
53

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54

(c) **FEEL** for the flow of air on your cheek (see Figure 2-6).

(d) **PERFORM** rescue breathing if the casualty does not resume breathing spontaneously.

NOTE

If the casualty resumes breathing, monitor and maintain the open airway. He should be transported to an MTF, as soon as practical.

2-8. Perform Rescue Breathing

(1) Use the thumb and index finger of the hand on the patient's forehead to pinch off his nostrils so that air will not escape.

(2) Take a deep breath, open your mouth very wide and place it around the outside of the patient's mouth making a seal.

(3) Blow air into the patient's mouth and at the same time look out of the corner of your eye to see if his chest is rising. If it is, the lungs are being ventilated. Ventilation should only be forceful enough to raise the patient's chest.

(4) Initially give four quick full breaths without allowing time for full lung deflation between breaths. If breathing has stopped, even for a short time, some of the small air sacs of the lungs collapse. Four initial breaths maintain positive pressure in the lungs, thereby, more effectively filling and ventilating the air sacs (Figure).

(5) If pulse is present but patient is not breathing you must perform rescue breathing. One breath is given every 5 seconds (for a respiratory rate of 12/minute). Between breaths you must put your head close to the patient's face to look, listen, and feel for spontaneous respirations. The pulse is checked once every minute.

(a) Establish unresponsiveness. Call for help. Turn or position the casualty.

(b) Open the airway.

(c) Check for breathing by placing your ear over the casualty's mouth and nose, and looking toward his chest.

1. **LOOK** for rise and fall of the casualty's chest (Figure 2-6)

2. **LISTEN** for sounds of breathing.

3. **FEEL** for breath on the side of your face. If the chest does not rise and fall and no air is exhaled, then, the casualty is not breathing.

4. **PERFORM** rescue breathing if the casualty is not breathing.

NOTE

Although the rescuer may notice that the casualty is making respiratory efforts, the airway may still be obstructed and opening the airway may

1 be all that is needed. If the casualty resumes breathing, the rescuer should
2 continue to maintain an open airway.
3

4
5 *Figure 2-6. Check for breathing.*
6

7
8 (2) *Mouth-to-mouth method.* In this method of rescue breathing, you inflate
9 the casualty's lungs with air from your lungs. This can be accomplished by blowing air into the
10 person's mouth. The mouth-to-mouth rescue breathing method is performed as follows:
11

12 (a) If the casualty is not breathing, place your hand on his forehead,
13 and pinch his nostrils together with the thumb and index finger of this same hand. Let this same
14 hand exert pressure on his forehead to maintain the backward head tilt and maintain an open
15 airway. With your other hand, keep your fingertips on the bony part of the lower jaw near the
16 chin and lift (Figure 2-7).
17

18
19 *Figure 2-7. Head tilt/chin lift.*
20

21
22 **NOTE**

23 If you suspect the casualty has a neck injury and you are using the
24 jaw thrust technique, close the nostrils by placing your cheek tightly
25 against them.
26

27 (b) Take a deep breath and place your mouth (in an airtight seal)
28 around the casualty's mouth (Figure 2-8). (If the injured person is small, cover both his nose and
29 mouth with your mouth, sealing your lips against the skin of his face.)
30

31
32 *Figure 2-8. Rescue breathing.*
33

34
35 (c) Blow two full breaths into the casualty's mouth (1 to 1 1/2 seconds
36 per breath), taking a breath of fresh air each time before you blow. Watch out of the corner of
37 your eye for the casualty's chest to rise. If the chest rises, sufficient air is getting into the
38 casualty's lungs. Therefore, proceed as described in step (b). If the chest does not rise, do the
39 following (1, 2, and 3 below) and then attempt to ventilate again.
40

41 1. Take corrective action immediately by reestablishing the
42 airway. Make sure that air is not leaking from around your mouth or out of the casualty's pinched
43 nose.
44

45 2. Reattempt to ventilate.
46

47 3. If the chest still does not rise, take the necessary action to
48 open an obstructed airway (paragraph 2-10).
49

50 **NOTE**

51 If the initial attempt to ventilate the casualty is unsuccessful,
52 reposition the casualty's head and repeat rescue breathing. Improper
53 chin and head positioning is the most common cause of difficulty
54 with ventilation. If the casualty cannot be ventilated after repositioning

1 the head, proceed with foreign-body airway obstruction maneuvers (see
2 paragraph 2-10).

3
4 (d) After giving two breaths which cause the chest to rise, attempt to
5 locate a pulse on the casualty. Feel for a pulse on the side of the casualty's neck closest to you by
6 placing the first two fingers (index and middle fingers) of your hand on the groove beside the
7 casualty's Adam's apple (carotid pulse) (Figure 2-9). (Your thumb should not be used for pulse
8 taking because you may confuse your pulse beat with that of the casualty.) Maintain the airway
9 by keeping your other hand on the casualty's forehead. Continue checking for the pulse for 5 to
10 10 seconds to determine if there is one.

11
12
13 *Figure 2-9. Placement of fingers to detect pulse.*

14
15
16 1. If a pulse is found and the casualty is breathing-- **STOP**;
17 allow the casualty to breathe on his own. If possible, keep him warm and comfortable.

18
19 2. If a pulse is found and the casualty is not breathing,
20 continue rescue breathing.

21
22 3. If a pulse is not found, immediately seek medical personnel
23 for help.

24
25 *b. Mouth-to-Mouth Resuscitation.* Mouth-to-mouth (for mouth-to-nose
26 resuscitation, refer to paragraph *c* below) is performed at the rate of about one breath every 5
27 seconds (12 breaths per minute) with rechecks for pulse and breathing after every 12 breaths.
28 Rechecks can be accomplished in 3 to 5 seconds. See steps one through seven (below) for
29 specifics.

30
31 **NOTE**

32 Send another soldier, if available, to get medical assistance.
33 Do not stop rescue breathing to get help.

34
35 (1) If the casualty is not breathing, pinch his nostrils together with the thumb
36 and index finger of the hand on his forehead and let this same hand exert pressure on the
37 forehead to maintain the backward head tilt (Figure 2-7).

38
39 (2) Take a deep breath and place your mouth (in an airtight seal) around the
40 casualty's mouth (Figure 2-8).

41
42 (3) Blow a quick breath into the casualty's mouth forcefully to cause his chest
43 to rise. If the casualty's chest rises, sufficient air is getting into his lungs.

44
45 (4) When the casualty's chest rises, remove your mouth from his mouth and
46 listen for the return of air from his lungs (exhalation).

47
48 (5) Repeat this procedure (mouth-to-mouth resuscitation) at a rate of one
49 breath every 5 seconds to achieve 12 breaths per minute. Use the following count: "one, one-
50 thousand; two, one-thousand; three, one-thousand; four, one-thousand; **BREATH**; one, one-
51 thousand;" and so forth. To achieve a rate of one breath every 5 seconds, the breath must be
52 given on the *fifth* count.

1 (6) Feel for a pulse after every 12th breath. This check should take about 3 to
2 5 seconds. If a pulse beat is not found, seek medical personnel for help.
3

4 (7) Continue rescue breathing until the casualty starts to breathe on his own,
5 until another person relieves you, or until you are too tired to continue. Monitor pulse and return
6 of spontaneous breathing after every few minutes of rescue breathing. If spontaneous breathing
7 returns, monitor the casualty closely. The casualty should be transported to an MTF. Maintain
8 an open airway and be prepared to resume rescue breathing if necessary.
9

10 c. *Mouth-to-Nose Method.* Use this method if you cannot perform mouth-to-mouth
11 rescue breathing because the casualty has a severe jaw fracture or mouth wound or his jaws are
12 tightly closed by spasms. The mouth-to-nose method is performed in the same way as the
13 mouth-to-mouth method except that you blow into the nose while you hold the lips closed with
14 one hand at the chin. You then remove your mouth to allow the casualty to exhale passively. It
15 may be necessary to separate the casualty's lips to allow the air to escape during exhalation.
16
17

18 **2-9. Airway Obstructions**

19
20 In order for oxygen from the air to flow to and from the lungs, the upper airway must be
21 unobstructed.
22

23 a. Upper airway obstructions often occur because--

24
25 (1) The casualty's tongue may fall back into his throat while he is
26 unconscious. The tongue *falls back and obstructs* the airway, it *is not swallowed* by the casualty.
27

28 **NOTE**

29 Ensure the correct positioning and maintenance of the open airway
30 for an injured or unconscious casualty.
31

32 (2) Foreign bodies become lodged in the throat. These obstructions usually
33 occur while eating. Choking on food (usually meat) is associated with--
34

- 35 • Attempting to swallow large pieces of poorly chewed food.
- 36 • Drinking alcohol.
- 37 • Slipping dentures.

38
39
40
41 (3) The contents of the stomach are regurgitated and may block the airway.

42 (4) Blood clots may form as a result of head and facial injuries.
43
44

45 b. Some upper airway obstructions may be prevented by taking the following
46 precautions:
47

- 48 • Cutting food into small pieces and taking care to chew slowly and
49 thoroughly.
 - 50 • Avoiding laughing and talking when chewing and swallowing.
 - 51 • Restricting alcohol intake while eating meals.
- 52
53
54

1 c. Upper airway obstruction may cause either partial or complete airway blockage.

2
3 (1) *Partial airway obstruction.* The casualty may still have an air exchange.
4 A good air exchange means that the casualty can cough forcefully, though he may be wheezing
5 between coughs. You, the rescuer, should not interfere, and should encourage the casualty to
6 cough up the object obstructing his airway on his own. A poor air exchange may be indicated by
7 weak coughing with a high pitched noise between coughs. Further, the casualty may show signs
8 of shock (paragraph 1-6b[5]) indicating a need for oxygen. You should assist the casualty and
9 treat him as though he had a complete obstruction.

10
11 (2) *Complete airway obstruction.* A complete obstruction (no air exchange) is
12 indicated if the casualty cannot speak, breathe, or cough at all. He may be clutching his neck and
13 moving erratically. In an unconscious casualty, a complete obstruction is also indicated if after
14 opening his airway you cannot ventilate him.

15 16 17 **2-10. Opening the Obstructed Airway--Conscious Casualty**

18
19 Clearing a conscious casualty's airway obstruction can be performed with the casualty either
20 standing or sitting and by following a relatively simple procedure.

21 **WARNING**

22
23 Once an obstructed airway occurs, the brain will develop an
24 oxygen deficiency resulting in unconsciousness. Death will
25 follow rapidly if breathing is not promptly restored.

26
27 a. Ask the casualty if he can speak or if he is choking. Check for the universal
28 choking sign (Figure 2-10).

29
30
31 *Figure 2-10. Universal sign of choking.*

32
33
34 b. If the casualty can speak, encourage him to attempt to cough; the casualty still has
35 a good air exchange. If he is able to speak or cough effectively, **DO NOT** interfere with his
36 attempts to expel the obstruction.

37
38 c. Listen for high pitched sounds when the casualty breathes or coughs (poor air
39 exchange). If there is poor air exchange or no breathing, **CALL** for **HELP** and immediately
40 deliver manual thrusts (either an abdominal or chest thrust).

41 **NOTE**

42
43 The manual thrust with the hands centered between the waist and
44 the rib cage is called an abdominal thrust (or Heimlich maneuver).
45 The chest thrust (the hands are centered in the middle of the breastbone)
46 is used only for an individual in the advanced stages of pregnancy, in the
47 markedly obese casualty, or if there is a significant abdominal wound.

48
49 (1) Apply **ABDOMINAL THRUSTS**. This can be accomplished by using
50 the following procedures:

51
52 (a) Stand behind the casualty and wrap your arms around his waist.

1 (b) Make a fist with one hand and grasp it with the other. The thumb
2 side of your fist should be against the casualty's abdomen, in the midline and slightly above the
3 casualty's navel, but well below the tip of the breastbone (Figure 2-11).
4

5
6 *Figure 2-11. Anatomical view of abdominal thrust procedure.*
7

8
9 (c) Press the fists into the abdomen with a quick backward and upward
10 thrust (Figure 2-12).
11

12
13 *Figure 2-12. Profile view of abdominal thrust.*
14

15
16 (d) Each thrust should be a separate and distinct movement.
17

18 **NOTE**

19 Continue performing abdominal thrusts until the obstruction is
20 expelled or the casualty becomes unconscious.
21

22 (e) If the casualty becomes unconscious, call for help as you proceed
23 with steps to open the airway and perform rescue breathing. (Refer to paragraph 2-7b for
24 information on how to perform mouth-to-mouth resuscitation.)
25

26 (2) Apply **CHEST THRUSTS**. An alternate technique to the abdominal
27 thrust is the chest thrust. This technique is useful when the casualty has an abdominal wound,
28 when the casualty is pregnant, or when the casualty is so large that you cannot wrap your arms
29 around the abdomen. To apply chest thrusts with casualty sitting or standing.
30

31 (a) Stand behind the casualty and wrap your arms around his chest
32 with your arms under his armpits.
33

34 (b) Make a fist with one hand and place the thumb side of the fist in
35 the middle of the breastbone (take care to avoid the tip of the breastbone and the margins of the
36 ribs).
37

38 (c) Grasp the fist with the other hand and exert thrusts (Figure 2-13).
39
40

41 *Figure 2-13. Profile view of chest thrust.*
42

43
44 (d) Each thrust should be delivered slowly, distinctly, and with the
45 intent of relieving the obstruction.
46

47 (e) Perform chest thrusts until the obstruction is expelled or the
48 casualty becomes unconscious.
49

50 (f) If the casualty becomes unconscious, call for help as you proceed
51 with steps to open the airway and perform rescue breathing.
52
53

54 **2-11. Open an Obstructed Airway--Casualty Lying Down or Unconscious**

1
2 The following procedures are used to expel an airway obstruction in a casualty who is lying
3 down, who becomes unconscious, or who is found unconscious (the cause unknown):
4

5 • If a conscious casualty who is choking becomes unconscious, call for help, open
6 the airway, perform a finger sweep, and attempt rescue breathing (paragraphs 2-3 through 2-7).
7 If you still cannot administer rescue breathing due to an airway blockage, then remove the airway
8 obstruction using the procedures as in *b* below.
9

10 • If a casualty is unconscious when you find him (the cause unknown), assess or
11 evaluate the situation, call for help, position the casualty on his back, open the airway, establish
12 breathlessness, and attempt to perform rescue breathing (paragraphs 2-3 through 2-7).
13

14 *a.* Open the airway and attempt rescue breathing (refer to paragraph 2-7 for
15 information on how to perform mouth-to-mouth resuscitation).
16

17 *b.* If still unable to ventilate the casualty, perform 6 to 10 manual (abdominal or
18 chest) thrusts.
19

20 (1) To perform the abdominal thrusts:
21

22 *(a)* Kneel astride the casualty's thighs (Figure 2-14).
23
24

25 *Figure 2-14. Abdominal thrust on unconscious casualty.*
26
27

28 *(b)* Place the heel of one hand against the casualty's abdomen (in the
29 midline slightly above the navel but well below the tip of the breastbone). Place your other hand
30 on top of the first one. Point your fingers toward the casualty's head.
31

32 *(c)* Press into the casualty's abdomen with a quick, forward and
33 upward thrust. You can use your body weight to perform the maneuver. Deliver each thrust
34 slowly and distinctly.
35

36 *(d)* Repeat the sequence of abdominal thrusts, finger sweep, and rescue
37 breathing (attempt to ventilate) as long as necessary to remove the object from the obstructed
38 airway.
39

40 *(e)* If the casualty's chest rises, proceed to feeling for pulse.
41

42 (2) To perform chest thrusts:
43

44 *(a)* Place the unconscious casualty on his back, face up, and open his
45 mouth. Kneel close to the side of the casualty's body.
46

47 *1.* Locate the lower edge of the casualty's ribs with your
48 fingers. Run the fingers up along the rib cage to the notch (Figure 2-15A).
49

50 *2.* Place the middle finger on the notch and the index finger
51 next to the middle finger on the lower edge of the breastbone. Place the heel of the other hand on
52 the lower half of the breastbone next to the two fingers (Figure 2-15B).
53

1 3. Remove the fingers from the notch and place that hand on
2 top of the positioned hand on the breastbone, extending or interlocking the fingers (Figure 2-
3 15C).

4
5 4. Straighten and lock your elbows with your shoulders
6 directly above your hands without bending the elbows, rocking, or allowing the shoulders to sag.
7 Apply enough pressure to depress the breastbone 1 1/2 to 2 inches, then release the pressure
8 completely (Figure 2-15D). Do this 6 to 10 times. Each thrust should be delivered slowly and
9 distinctly. See Figure 2-16 for another view of the breastbone being depressed.

10
11
12 *Figure 2-15. Hand placement for chest thrust (Illustrated A-D).*

13
14
15 *Figure 2-16. Breastbone depressed 1 1/2 to 2 inches.*

16
17
18 (b) Repeat the sequence of chest thrust, finger sweep, and rescue
19 breathing as long as necessary to clear the object from the obstructed airway. See paragraph (3)
20 below.

21
22 (c) If the casualty's chest rises, proceed to feeling for his pulse.

23
24 (3) If you still cannot administer rescue breathing due to an airway
25 obstruction, then remove the airway obstruction using the procedures in steps (a) and (b) below.

26
27 (a) Place the casualty on his back, face up, turn the unconscious
28 casualty as a unit, and call out for help.

29
30 (b) Perform finger sweep, keep casualty face up, use tongue-jaw lift to
31 open mouth.

32
33 1. Open the casualty's mouth by grasping both his tongue and
34 lower jaw between your thumb and fingers and lifting (tongue-jaw lift) (Figure 2-17). If you are
35 unable to open his mouth, cross your fingers and thumb (crossed-finger method) and push his
36 teeth apart (Figure 2-18) by pressing your thumb against his upper teeth and pressing your finger
37 against his lower teeth.

38
39
40 *Figure 2-17. Opening casualty's mouth (tongue-jaw lift).*

41
42
43 *Figure 2-18. Opening casualty's mouth (crossed-finger method).*

44
45
46 2. Insert the index finger of the other hand down along the
47 inside of his cheek to the base of the tongue. Use a hooking motion from the side of the mouth
48 toward the center to dislodge the foreign body (Figure 2-19).

49
50
51 *Figure 2-19. Using finger to dislodge a foreign body.*

52
53
54 **WARNING**

1 Take care not to force the object deeper into the airway by pushing
2 it with the finger.
3
4
5

6 **Section II. STOP THE BLEEDING AND PROTECT THE WOUND**

7
8

9 **2-12. General**

10

11 The longer a soldier bleeds from a major wound, the less likely he will be able to survive his
12 injuries. It is, therefore, important that the first aid provider promptly stop the external bleeding.
13
14

15 **2-13. Clothing**

16

17 In evaluating the casualty for location, type, and size of the wound or injury, cut or tear his
18 clothing and carefully expose the entire area of the wound. This procedure is necessary to avoid
19 further contamination. Clothing stuck to the wound should be left in place to avoid further
20 injury. DO NOT touch the wound; keep it as clean as possible.
21

22 **WARNING**

23 DO NOT REMOVE protective clothing in a chemical environment.
24 Apply dressings over the protective clothing.
25
26

27 **2-14. Entrance and Exit Wounds**

28

29 Before applying the dressing, carefully examine the casualty to determine if there is more than
30 one wound. A missile may have entered at one point and exited at another point. The *EXIT*
31 wound is usually *LARGER* than the entrance wound.
32
33

34 **WARNING**

35 The casualty should be continually monitored for development of
36 conditions which may require the performance of necessary basic
37 lifesaving measures, such as clearing the airway and mouth-to-mouth
38 resuscitation. All open (or penetrating) wounds should be checked for
39 a point of entry and exit and first aid measures applied accordingly.
40

41 **WARNING**

42 If the missile lodges in the body (fails to exit), DO NOT attempt to
43 remove it or probe the wound. Apply a dressing. If there is an object
44 extending from (impaled in) the wound, DO NOT remove the object.
45 Apply a dressing around the object and use additional improvised bulky
46 materials/dressings (use the cleanest material available) to build up the
47 area around the object. Apply a supporting bandage over the bulky
48 materials to hold them in place.
49

50 **2-15. Field Dressing**

51

52 *a.* Use the casualty's field dressing; remove it from the wrapper and grasp the tails of
53 the dressing with both hands (Figure 2-20).
54

1
2 *Figure 2-20. Grasping tails of dressing with both hands.*
3

4
5 **WARNING**

6 DO NOT touch the white (sterile) side of the dressing, and DO
7 NOT allow it to come in contact with any surface other than
8 the wound.
9

10 *b.* Hold the dressing directly over the wound with the white side down. Pull the
11 dressing open (Figure 2-21) and place it directly over the wound (Figure 2-22).
12

13
14 *Figure 2-21. Pulling dressing open.*
15

16
17 *Figure 2-22. Placing dressing directly on wound.*
18

19
20 *c.* Hold the dressing in place with one hand. Use the other hand to wrap one of the
21 tails around the injured part, covering about one-half of the dressing (Figure 2-23). Leave
22 enough of the tail for a knot. If the casualty is able, he may assist by holding the dressing in
23 place.
24

25
26 *Figure 2-23. Wrapping tail of dressing around injured part.*
27

28
29 *d.* Wrap the other tail in the opposite direction until the remainder of the dressing is
30 covered. The tails should seal the sides of the dressing to keep foreign material from getting
31 under it.
32

33 *e.* Tie the tails into a nonslip knot over the outer edge of the dressing (Figure 2-24).
34 **DO NOT TIE THE KNOT OVER THE WOUND.** In order to allow blood to flow to the rest
35 of an injured limb, tie the dressing firmly enough to prevent it from slipping but without causing
36 a tourniquet-like effect; that is, the skin beyond the injury becomes cool, blue, or numb.
37

38
39 *Figure 2-24. Tails tied into nonslip knot.*
40

41
42 **2-16. Manual Pressure**
43

44 *a.* If bleeding continues after applying the sterile field dressing, direct manual
45 pressure may be used to help control bleeding. Apply such pressure by placing a hand on the
46 dressing and exerting firm pressure for 5 to 10 minutes (Figure 2-25). The casualty may be asked
47 to do this himself if he is conscious and can follow instructions.
48

49
50 *Figure 2-25. Direct manual pressure applied.*
51

52
53 *b.* Elevate an injured limb slightly above the level of the heart to reduce the bleeding
54 (Figure 2-26).

1
2
3 *Figure 2-26. Injured limb elevated.*

4
5 **WARNING**

6 DO NOT elevate a suspected fractured limb unless it has been
7 properly splinted.

8
9 c. If the bleeding stops, check shock; administer first aid for shock as necessary. If
10 the bleeding continues, apply a pressure dressing.

11
12
13 **2-17. Pressure Dressing**

14
15 Pressure dressings aid in blood clotting and compress the open blood vessel. If bleeding
16 continues after the application of a field dressing, manual pressure, and elevation, then a pressure
17 dressing must be applied as follows:

18
19 a. Place a wad of padding on top of the field dressing, directly over the wound
20 (Figure 2-27). Keep the injured extremity elevated.

21
22
23 *Figure 2-27. Wad of padding on top of field dressing.*

24
25
26 **NOTE**

27 Improvised bandages may be made from strips of cloth. These
28 strips may be made from T-shirts, socks, or other garments.

29
30 b. Place an improvised dressing (or cravat, if available) over the wad of padding
31 (Figure 2-28). Wrap the ends tightly around the injured limb, covering the previously placed
32 field dressing (Figure 2-29).

33
34
35 *Figure 2-28. Improvised dressing over wad of padding.*

36
37
38 *Figure 2-29. Ends of improvised dressing wrapped tightly around limb.*

39
40
41 c. Tie the ends together in a nonslip knot, directly over the wound site (Figure 2-30).
42 DO NOT tie so tightly that it has a tourniquet-like effect. If bleeding continues and all other
43 measures have failed, or if the limb is severed, then apply a tourniquet. Use the tourniquet as a
44 **LAST RESORT**. When the bleeding stops, check for shock; administer first aid for shock as
45 necessary.

46
47
48 *Figure 2-30. Ends of improvised dressing tied together in nonslip knot.*

49
50 **NOTE**

51 Wounded extremities should be checked periodically for adequate
52 circulation. The dressing must be loosened if the extremity becomes
53 cool, blue, or numb.

NOTE

If bleeding continues and all other measures have failed (dressings and covering wound, applying direct manual pressure, elevating the limb above the heart level, and applying a pressure dressing while maintaining limb elevation) *then apply digital pressure*. (Refer to Appendix C for pressure points.)

2-18. Tourniquet

A soldier whose arm or leg has been completely amputated may not be bleeding when first discovered, but a tourniquet should be applied anyway. This absence of bleeding is due to the body's normal defenses (contraction of blood vessels) as a result of the amputation, but after a period of time bleeding will start as the blood vessels relax. Bleeding from a major artery of the thigh, lower leg, or arm and bleeding from multiple arteries (which occurs in a traumatic amputation) may prove to be beyond control by manual pressure. If the pressure dressing under firm hand pressure becomes soaked with blood and the wound continues to bleed, apply a tourniquet. A tourniquet is a constricting band placed around an arm or leg to control bleeding.

WARNING

Casualty should be continually monitored for development of conditions which may require the performance of necessary basic lifesaving measures, such as: clearing the airway, performing mouth-to-mouth resuscitation, preventing shock, and/or bleeding control. All open (or penetrating) wounds should be checked for a point of entry or exit and treated accordingly.

The tourniquet should not be used unless a pressure dressing has failed to stop the bleeding or an arm or leg has been cut off. On occasion, tourniquets have injured blood vessels and nerves. If left in place too long, a tourniquet can cause loss of an arm or leg. Once applied, it must stay in place, and the casualty must be taken to the nearest MTF as soon as possible. **DO NOT loosen or release a tourniquet after it has been applied and the bleeding has stopped.**

a. Improvising a Tourniquet. In the absence of a specially designed tourniquet, a tourniquet may be made from a strong, pliable material, such as gauze or muslin bandages, clothing, or kerchiefs. An improvised tourniquet is used with a rigid stick-like object. To minimize skin damage, ensure that the improvised tourniquet is at least 2 inches wide.

WARNING

The tourniquet must be easily identified or easily seen.

WARNING

DO NOT use wire or shoestring for a tourniquet band.

WARNING

A tourniquet is only used on arm(s) or leg(s) where there is danger of the casualty losing his life.

b. Placing the Improvised Tourniquet.

(1) Place the tourniquet around the limb, between the wound and the body trunk (or between the wound and the heart). Place the tourniquet 2 to 4 inches from the edge of the wound site (Figure 2-31). Never place it directly over a wound or fracture or directly on a

1 joint (wrist, elbow, or knee). For wounds just below a joint, place the tourniquet just above and
2 as close to the joint as possible.
3

4
5 *Figure 2-31. Tourniquet 2 to 4 inches above wound.*
6

7
8 (2) The tourniquet should have padding underneath. If possible, place the
9 tourniquet over the smoothed sleeve or trouser leg to prevent the skin from being pinched or
10 twisted. If the tourniquet is long enough, wrap it around the limb several times, keeping the
11 material as flat as possible. Damaging the skin may deprive the surgeon of skin required to cover
12 an amputation. Protection of the skin also reduces pain.
13

14 c. *Applying the Tourniquet.*

15
16 (1) Tie a half-knot. (A half-knot is the same as the first part of tying a shoe
17 lace.)

18
19 (2) Place a stick (or similar rigid object) on top of the half-knot (Figure 2-32).
20

21
22 *Figure 2-32. Rigid object on top of half-knot.*
23

24
25 (3) Tie a full knot over the stick (Figure 2-33).
26

27
28 *Figure 2-33. Full knot over rigid object.*
29

30 (4) Twist the stick (Figure 2-34) until the tourniquet is tight around the limb
31 and/or the bright red bleeding has stopped. In the case of amputation, dark oozing blood may
32 continue for a short time. This is the blood trapped in the area between the wound and
33 tourniquet.
34

35
36 *Figure 2-34. Stick twisted.*
37

38
39 (5) Fasten the tourniquet to the limb by looping the free ends of the tourniquet
40 over the ends of the stick. Then bring the ends around the limb to prevent the stick from
41 loosening. Tie them together on the side of the limb (Figure 2-35A and B).
42

43
44 *Figure 2-35. Free ends looped (Illustrated A and B).*
45

46
47 **NOTE**

48 Other methods of securing the stick may be used as long as the
49 stick does not unwind and no further injury results.
50

51 **NOTE**

52 If possible, save and transport any severed (amputated) limbs or
53 body parts with (but out of sight of) the casualty.
54

1 (6) DO NOT cover the tourniquet--you should leave it in full view. If the
2 limb is missing (total amputation), apply a dressing to the stump.
3

4 (7) Mark the casualty's forehead, if possible, with a "T" to indicate a
5 tourniquet has been applied. If necessary, use the casualty's blood to make this mark.
6

7 (8) Check and treat for shock.
8

9 (9) Seek medical aid.
10

11 CAUTION

12 Do not loosen or release the tourniquet once it has been applied
13 because it could enhance the probability of shock.
14

15 Section III. CHECK FOR SHOCK AND ADMINISTER FIRST AID MEASURES 16

17 2-19. General 18

19 The term *shock* has a variety of meanings. In medicine, it refers to a collapse of the body's
20 cardiovascular system which includes an inadequate supply of blood to the body's tissues. Shock
21 stuns and weakens the body. When the normal blood flow in the body is upset, death can result.
22 Early recognition and proper first aid may save the casualty's life.
23
24

25 2-20. Causes and Effects 26

27 *a.* There are three basic mechanisms associated with shock. These are--
28
29 • The heart is damaged and fails to work as a pump.
30
31 • Blood loss (heavy bleeding) causes the volume of fluid within the vascular
32 system to be insufficient.
33
34 • The blood vessels dilate (open wider) so that the blood within the system
35 (even though it is a normal volume [the casualty is not bleeding or dehydrated]) is insufficient to
36 provide adequate circulation within the body.
37
38

39 *b.* Shock may be the result of a number of conditions. These include--
40
41 • Dehydration.
42
43 • Allergic reaction to foods, drugs, insect stings, and snakebites.
44
45 • Significant loss of blood.
46
47 • Reaction to the sight of a wound, blood, or other traumatic scene.
48
49 • Traumatic injuries, such as--
50
51 • Burns.
52
53 • Gunshot or shrapnel wounds.
54

- 1
- 2 • Crush injuries.
- 3
- 4 • Blows to the body (which can cause broken bones or damage to
- 5 internal organs).
- 6
- 7 • Head injuries.
- 8
- 9 • Penetrating wounds (such as from a knife, bayonet, or missile).
- 10

11 **2-21. Signs and Symptoms of Shock**

12
13 Examine the casualty to see if he has any of the following signs and symptoms:

- 14 • Sweaty but cool skin (clammy skin).
- 15
- 16 • Paleness of skin (in dark-skinned individuals they may have a grayish look to their
- 17 skin).
- 18
- 19 • Restlessness, nervousness.
- 20
- 21 • Thirst.
- 22
- 23 • Loss of blood (bleeding).
- 24
- 25 • Confusion (or loss of awareness).
- 26
- 27 • Faster-than-normal breathing rate.
- 28
- 29 • Blotchy or bluish skin (especially around the mouth and lips).
- 30
- 31 • Nausea and/or vomiting.
- 32
- 33
- 34

35 **2-22. First Aid Measures for Shock**

36
37 In the field, the first aid procedures administered for shock are identical to procedures that would
38 be performed *to prevent shock*. When treating a casualty, assume that shock is present or will
39 occur shortly. By waiting until actual signs and symptoms of shock are noticeable, the rescuer
40 may jeopardize the casualty's life.

41
42 *a. Position the Casualty.* (DO NOT move the casualty or his limbs if suspected
43 fractures have not been splinted. See Chapter 4 for details.)

- 44
- 45 (1) Move the casualty to cover, if cover is available and the situation permits.
- 46
- 47 (2) Lay the casualty on his back.
- 48

49 **NOTE**

50 A casualty in shock from a chest wound or one who is experiencing
51 breathing difficulty, may breathe easier in a sitting position. If this
52 is the case, allow him to sit upright, but monitor carefully in case his
53 condition worsens.

54

1 (3) Elevate the casualty's feet higher than the level of his heart. Use a stable
2 object (field pack or rolled up clothing) so that his feet will not slip off (Figure 2-36).
3

4 **WARNING**

5 DO NOT elevate legs if the casualty has an unsplinted broken leg,
6 head injury, or abdominal injury.
7

8
9 *Figure 2-36. Clothing loosened and feet elevated.*
10

11 **WARNING**

12 Check casualty for leg fracture(s) and splint, if necessary, before
13 elevating his feet. For a casualty with an abdominal wound, place
14 his knees in an upright (flexed) position.
15

16
17 (4) Loosen clothing at the neck, waist, or wherever it may be binding.
18

19 **CAUTION**

20 DO NOT loosen or remove protective clothing in a chemical
21 environment.
22

23 (5) Prevent chilling or overheating. The key is to maintain body temperature.
24 In cold weather, place a blanket or other like item over him to keep him warm and under him to
25 prevent chilling (Figure 2-37). However, if a tourniquet has been applied, leave it exposed (if
26 possible). In hot weather, place the casualty in the shade and protect him from becoming chilled;
27 however, avoid the excessive use of blankets or other coverings.
28

29
30 *Figure 2-37. Body temperature maintained.*
31

32
33 (6) Calm the casualty. Throughout the entire procedure of providing first aid
34 for a casualty, the rescuer should reassure the casualty and keep him calm. This can be done by
35 being authoritative (taking charge) and by showing self-confidence. Assure the casualty that you
36 are there to help him.
37

38 (7) Seek medical aid.
39

40 *b. Food and/or Drink.* When providing first aid for shock, DO NOT give the
41 casualty any food or drink. If you must leave the casualty or if he is unconscious, turn his head to
42 the side to prevent him from choking should he vomit (Figure 2-38).
43

44
45 *Figure 2-38. Casualty's head turned to side.*
46

47
48 *c. Evaluate Casualty.* Continue to evaluate the casualty until medical personnel
49 arrive or the casualty is transported to an MTF.

CHAPTER 3

FIRST AID FOR SPECIFIC INJURIES

3-1. General

Basic lifesaving steps are discussed in Chapters 1 and 2; they apply to first aid measures for all injuries. Certain types of wounds and burns will require special precautions and procedures when applying these measures. This chapter discusses first aid procedures for special wounds of the head, face, and neck; chest and stomach wounds; and burns. It also discusses the techniques for applying dressings and bandages to specific parts of the body.

3-2. Head, Neck, and Facial Injuries

a. Head Injuries.

(1) Head injuries range from minor abrasions or cuts on the scalp to severe brain injuries that may result in unconsciousness and sometimes death. Head injuries are also classified as open or closed wounds. An open wound is one that is visible, has a break in the skin, and usually has evidence of bleeding. A closed wound may be visible (such as a depression in the skull) or the first aid provider may not be able to see any apparent injury (such as internal bleeding). Some head injuries result in unconsciousness; however, a soldier may have a serious head wound and still be conscious. Due to the anatomical structure of the head and neck and the potential for worsening a head or spinal injury, casualties with head and neck injuries should be treated as though they also have a spinal injury. Unless the casualty's life is in immediate danger (such as being close to a burning vehicle), all precautions for stabilizing the head and neck prior to moving the casualty should be taken.

(2) Prompt first aid measures should be initiated for casualties with suspected head and neck injuries. The casualty may be conscious or unconscious. The conscious casualty may be able to provide information on the extent of his injuries or, as a result of the head injury, his thinking processes may not be working correctly. The signs and symptoms a first aid provider might observe are—

- Nausea and vomiting.
- Convulsions or twitches.
- Slurred speech.
- Confusion and loss of memory. (Does he know who he is? Does he know where he is? Does he know what day it is?)

- 1 • Recent unconsciousness
- 2
- 3 • Dizziness.
- 4
- 5 • Sleepiness.
- 6
- 7 • Blurred vision, unequal pupils, or bruising (black eyes).
- 8
- 9 • Paralysis (partial or full).
- 10
- 11 • Complaint of headache.
- 12
- 13 • Bleeding from the scalp, nose, or ears.
- 14
- 15 • Deformity of the head (depression or swelling).
- 16

17 *b. Neck Injuries.* Neck injuries may result in heavy bleeding. Apply pressure above
18 and below the injury, *but do not interfere with the breathing process*, and attempt to control the
19 bleeding. Apply a dressing. Always evaluate the casualty for a possible neck fracture/spinal cord
20 injury; if suspected, seek medical treatment immediately.

21 **NOTE**

22 Establish and maintain the airway in cases of facial or neck
23 injuries. If a neck fracture or spinal cord injury is suspected,
24 immobilize the injury and, if necessary, perform basic life
25 support measures.

26

27

28 *c. Facial Injuries.* Soft tissue injuries of the face and scalp are common. Abrasions
29 (scrapes) of the skin cause no serious problems. Contusions (injury without a break in the skin)
30 usually cause swelling. A contusion of the scalp looks and feels like a lump. Laceration (cut)
31 and avulsion (torn away tissue) injuries are also common. Avulsions are frequently caused when
32 a sharp blow separates the scalp from the skull beneath it. Because the face and scalp are richly
33 supplied with blood vessels (arteries and veins), wounds of these areas usually bleed heavily.

34 **3-3. General First Aid Measures**

35

36

37

38 *a. General Considerations.* The casualty with a head injury (or suspected head
39 injury) should be continually monitored for the development of conditions that *may require* the
40 performance of the necessary basic lifesaving measures. After initiating first aid measures
41 request medical assistance and evacuation. If dedicated medical evacuation assets are not
42 available, transport the casualty to an MTF as soon as the situation permits. The first aid
43 provider should not attempt to remove a protruding object from the head or give the casualty
44 anything to eat or drink. Further the first aid provider should be prepared to—

45

- 1 • Clear the airway.
- 2
- 3 • Control bleeding (external).
- 4
- 5 • Administer first aid measures for shock.
- 6
- 7 • Keep the casualty warm.
- 8
- 9 • Protect the wound.

10
11 *b. Unconscious Casualty.* An unconscious casualty does not have control of all of
12 his body's functions and may choke on his tongue, blood, vomitus, or other substances.

13
14 (1) *Breathing.* The brain requires a constant supply of oxygen. A bluish (or
15 in an individual with dark skin--grayish) color of skin around the lips and nail beds indicates that
16 the casualty is not receiving enough air (oxygen). Immediate action must be taken to clear the
17 airway, to position the casualty on his side, or to initiate rescue breathing.

18
19 (2) *Bleeding.* Bleeding from a head injury usually comes from blood vessels
20 within the scalp. Bleeding can also develop inside the skull or within the brain. In most
21 instances visible bleeding from the head can be controlled by proper application of the field first
22 aid dressing.

23
24 **CAUTION**

- 25 • DO NOT attempt to put unnecessary pressure on the wound
- 26 or attempt to push any brain matter back into the head (skull).
- 27 • DO NOT apply a pressure dressing.

28
29 *c. Concussion.* If an individual receives a heavy blow to the head or face, he may
30 suffer a brain concussion, which is an injury to the brain that involves a temporary *loss of some*
31 *or all* of the brain's ability to function. For example, the casualty may not breathe properly for a
32 short period of time, or he may become confused and stagger when he attempts to walk. A
33 concussion may only last for a short period of time. However, if a casualty is suspected of
34 having suffered a concussion, he should be transported to an MTF as soon as conditions permit.

35
36 *d. Convulsions.* Convulsions (seizures/involuntary jerking) may occur even after a
37 mild head injury. When a casualty is convulsing, protect him from hurting himself. Take the
38 following measures:

- 39
- 40 (1) Ease him to the ground if he is standing or sitting.
- 41
- 42 (2) Support his head and neck.
- 43
- 44 (3) Maintain his airway.
- 45

- 1 (4) Protect him from further injury (such as hitting close-by objects).
2

3 **NOTE**

- 4 • DO NOT forcefully hold the arms and legs if they are jerking
5 because this can lead to broken bones.
6 • DO NOT force anything between the casualty's teeth--especially
7 if they are tightly clenched because this may obstruct the casualty's
8 airway.
9 • Maintain the casualty's airway if necessary.

10
11 e. *Brain Damage.* In severe head injuries where brain tissue is protruding, *leave the*
12 *wound alone*; carefully place a first aid dressing over the tissue to protect it from further
13 contamination. *DO NOT remove or disturb any foreign matter that may be in the wound.*
14 Position the casualty so that his head is higher than his body. Keep him warm and *seek medical*
15 *aid immediately.*

16
17 **NOTE**

18 If there is an object extending from the wound, DO NOT remove
19 the object. Improvise bulky dressings from the cleanest material
20 available and place this material around the protruding object for
21 support, then apply the field dressing.
22
23

24 **3-4. Dressings and Bandages**

25
26 a. *Head Wounds.*

- 27
28 (1) *Position the Casualty.*
29

30 **WARNING**

31 DO NOT move the casualty if you suspect he has sustained a neck,
32 spine or head injury (which produces any signs or symptoms other
33 than minor bleeding).
34

- 35 • If the casualty has a minor (superficial) scalp wound and is
36 conscious:
37
38 • Have the casualty sit up (unless other injuries prohibit or he
39 is unable to); OR
40
41 • If the casualty is lying down and is not accumulating fluids
42 or drainage in his throat, elevate his head slightly; OR
43

1 • If the casualty is bleeding from or into his mouth or throat,
2 turn his head to the side or position him on his side so that the airway will be clear. Avoid
3 putting pressure on the wound and place him on his uninjured side (Figure 3-1).
4

5
6 *Figure 3-1. Casualty lying on side opposite injury.*
7

8
9 • If the casualty is unconscious or has a severe head injury, then
10 suspect and treat him as having a potential neck or spinal injury, *immobilize and DO NOT move*
11 *the casualty.*
12

13 **NOTE**

14 If the casualty is choking and/or vomiting or is bleeding from or
15 into his mouth (thus compromising his airway), position him on
16 his uninjured side so that his airway will be clear.
17

18 **WARNING**

19 If it is necessary to turn a casualty with a suspected neck/spine
20 injury; roll the casualty gently onto his side, keeping the head,
21 neck, and body aligned while providing support for the head and
22 neck. **DO NOT** roll the casualty by yourself but seek assistance.
23 *Move him only if absolutely necessary*, otherwise keep the casualty
24 immobilized to prevent further damage to the neck/spine.
25

26 (2) *Expose the wound.* Remove the casualty's helmet (if necessary). In an
27 nuclear, biological, and chemical (NBC) environment, the first aid provider must leave the
28 casualty as much protection (such as protective mask, mission-oriented protective posture
29 (MOPP) overgarments) as possible. What items of protective equipment can be removed is
30 dependent upon the casualty's injuries (where on the body and what type), the MOPP level,
31 integrity of protective equipment (such as tears in the garment or mask seal), availability of
32 chemical protective shelters, and the tactical situation.
33

34 **WARNING**

35 **DO NOT** attempt to clean the wound or remove a protruding object.
36

37 **NOTE**

38 Always use the casualty's field dressing, not your own.
39

40 (3) *Apply a Dressing to a Wound of the Forehead or Back of Head.* To apply
41 a dressing to a wound of the forehead or back of the head—
42

43 (a) Remove the dressing from the wrapper.
44

45 (b) Grasp the tails of the dressing in both hands.

1
2 (c) Hold the dressing (white side down) directly over the wound. DO
3 NOT touch the white (sterile) side of the dressing or allow anything except the wound to come in
4 contact with it.

5
6 (d) Place it directly over the wound.

7
8 (e) Hold it in place with one hand. If the casualty is able, he may
9 assist.

10
11 (f) Wrap the first tail horizontally around the head; ensure the tail
12 covers the dressing (Figure 3-2).

13
14
15 *Figure 3-2. First tail of dressing wrapped horizontally around head.*

16
17
18 (g) Hold the first tail in place and wrap the second tail in the opposite
19 direction, covering the dressing (Figure 3-3).

20
21
22 *Figure 3-3. Second tail wrapped in opposite direction.*

23
24
25 (h) Tie a nonslip knot and secure the tails at the side of the head,
26 making sure they DO NOT cover the eyes or ears (Figure 3-4).

27
28
29 *Figure 3-4. Tails tied in nonslip knot at side of head.*

30
31
32 (4) *Apply a Dressing to a Wound on Top of the Head.* To apply a dressing to
33 a wound on top of the head—

34
35 (a) Remove the dressing from the wrapper.

36
37 (b) Grasp the tails of the dressing in both hands.

38
39 (c) Hold it (white side down) directly over the wound.

40
41 (d) Place it over the wound (Figure 3-5).

42
43
44 *Figure 3-5. Dressing placed over wound.*

1 (e) Hold it in place with one hand. If the casualty is able, he may
2 assist.

3
4 (f) Wrap one tail down under the chin (Figure 3-6), up in front of the
5 ear, over the dressing, and in front of the other ear.

6
7
8 *Figure 3-6. One tail of dressing wrapped under chin.*

9
10
11 **WARNING**

12 Ensure the tails remain wide and close to the front of the chin to
13 avoid choking the casualty.

14
15 (g) Wrap the remaining tail under the chin in the opposite direction
16 and up the side of the face to meet the first tail (Figure 3-7).

17
18
19 *Figure 3-7. Remaining tail wrapped under chin in opposite direction.*

20
21
22 (h) Cross the tails (Figure 3-8), bringing one around the forehead
23 (above the eyebrows) and the other around the back of the head (at the base of the skull) to a
24 point just above and in front of the opposite ear, and tie them using a nonslip knot (Figure 3-9).

25
26
27 *Figure 3-8. Tails of dressing crossed with one around forehead.*

28
29
30 *Figure 3-9. Tails tied in nonslip knot (in front of and above ear).*

31
32
33 (5) *Apply a Triangular Bandage to the Head.* To apply a triangular bandage
34 to the head—

35
36 (a) Turn the base (longest side) of the bandage up and center its base
37 on the center of the forehead, letting the point (apex) fall on the back of the neck (Figure 3-10A).

38
39 (b) Take the ends behind the head and cross the ends over the apex.

40
41 (c) Take them over the forehead and tie them (Figure 3-10B).

42
43 (d) Tuck the apex behind the crossed part of the bandage and/or secure
44 it with a safety pin, if available (Figure 3-10C).

45

1
2 *Figure 3-10. Triangular bandage applied to head (Illustrated A through C)*
3
4

5 (6) *Apply a Cravat Bandage to the Head.* To apply a cravat bandage to the
6 head—

7
8 (a) Place the middle of the bandage over the dressing (Figure 3-11A).
9

10 (b) Cross the two ends of the bandage in opposite directions
11 completely around the head (Figure 3-11B).
12

13 (c) Tie the ends over the dressing (Figure 3-11C).
14
15

16 *Figure 3-11. Cravat bandage applied to head (Illustrated A through C).*
17
18

19 *b. Eye Injuries.* The eye is a vital sensory organ, and blindness is a severe physical
20 handicap. Timely first aid of the eye may relieve pain and may also help to prevent shock,
21 permanent eye injury, and possible loss of vision. Because the eye is very sensitive, any injury
22 can be easily aggravated if it is improperly handled. Injuries of the eye may be quite severe.
23 Cuts of the eyelids can appear to be very serious, but if the eyeball is not involved, a person's
24 vision usually will not be damaged. However, lacerations (cuts) of the eyeball can cause
25 permanent damage or loss of sight.
26

27 (1) *Lacerated/torn eyelids.* Lacerated eyelids may bleed heavily, but bleeding
28 usually stops quickly. Cover the injured eye with a sterile dressing. DO NOT put pressure on
29 the wound because you may injure the eyeball. Handle torn eyelids very carefully to prevent
30 further injury. Place any detached pieces of the eyelid on a clean bandage or dressing and
31 immediately send them with the casualty to the medical facility.
32

33 (2) *Lacerated eyeball (injury to the globe).* Lacerations or cuts to the eyeball
34 may cause serious and permanent eye damage. Cover the injury with a loose sterile dressing.
35 DO NOT put pressure on the eyeball because additional damage may occur. An important point
36 to remember is that when one eyeball is injured, you should immobilize both eyes. This is done
37 by applying a bandage to both eyes. Because the eyes move together, covering both will lessen
38 the chances of further damage to the injured eye.
39

40 **CAUTION**

41 DO NOT apply pressure when there is a possible laceration of the
42 eyeball. The eyeball contains fluid. Pressure applied over the eye
43 will force the fluid out, resulting in permanent injury. APPLY
44 PROTECTIVE DRESSING WITHOUT ADDED PRESSURE.
45

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45

(1) Remove the dressing from its wrapper.

(2) Grasp the tails in both hands.

(3) Hold the dressing directly over the wound with the white side down and place it directly on the wound (Figure 3-15A).

(4) Hold the dressing in place with one hand (the casualty may assist if able). Wrap the top tail over the top of the head and bring it down in front of the ear (on the side opposite the wound), under the chin (Figure 3-15B) and up over the dressing to a point just above the ear (on the wound side).

Figure 3-14. Side of head or cheek wound.

Figure 3-15. Dressing placed directly on wound. Top tail wrapped over top of head, down in front of ear, and under chin (Illustrated A and B).

NOTE

When possible, avoid covering the casualty's ear with the dressing, as this will decrease his ability to hear.

(5) Bring the second tail under the chin, up in front of the ear (on the side opposite the wound), and over the head to meet the other tail (on the wound side) (Figure 3-16).

Figure 3-16. Bringing second tail under the chin.

(6) Cross the two tails (on the wound side) (Figure 3-17) and bring one end across the forehead (above the eyebrows) to a point just in front of the opposite ear (on the uninjured side).

Figure 3-17. Crossing the tails on the side of the wound.

(7) Wrap the other tail around the back of the head (at the base of the skull), and tie the two ends just in front of the ear on the uninjured side with a nonslip knot (Figure 3-18).

1 *Figure 3-18. Tying the tails of the dressing in a nonslip knot.*

2
3
4 *d. Ear Injuries.* Lacerated (cut) or avulsed (torn) ear tissue may not, in itself, be a
5 serious injury. Bleeding, or the drainage of fluids from the ear canal, however, may be a sign of
6 a head injury, such as a skull fracture. DO NOT attempt to stop the flow from the inner ear canal
7 nor put anything into the ear canal to block it. Instead, you should cover the ear lightly with a
8 dressing. For minor cuts or wounds to the external ear, apply a cravat bandage as follows:

9
10 (1) Place the middle of the bandage over the ear (Figure 3-19A).

11
12 (2) Cross the ends, wrap them in opposite directions around the head, and tie
13 them (Figures 3-19B and 3-19C).

14
15
16 *Figure 3-19. Applying cravat bandage to ear (Illustrated A through C).*

17
18
19 (3) If possible, place some dressing material between the back of the ear and
20 the side of the head to avoid crushing the ear against the head with the bandage.

21
22 *e. Nose Injuries.* Nose injuries generally produce bleeding. The bleeding may be
23 controlled by placing an ice pack over the nose, or pinching the nostrils together. The bleeding
24 may also be controlled by placing torn gauze (rolled) between the upper teeth and the lip.

25
26 **CAUTION**

27 DO NOT attempt to remove objects inhaled into the nose. An
28 untrained person who removes such an object could worsen the
29 casualty's condition and cause permanent injury.

30
31 *f. Jaw Injuries.* Before applying a bandage to a casualty's jaw, remove all foreign
32 material from the casualty's mouth. If the casualty is unconscious, check for obstructions in the
33 airway. When applying the bandage, allow the jaw enough freedom to permit passage of air and
34 drainage from the mouth.

35
36 (1) *Apply bandages attached to field first aid dressing to the jaw.* After
37 dressing the wound, apply the bandages using the same technique illustrated in Figures 3-5
38 through 3-8.

39
40 **NOTE**

41 The dressing and bandaging procedure outlined for the jaw serves
42 a twofold purpose. In addition to stopping the bleeding and protecting
43 the wound, it also immobilizes a fractured jaw.

44
45 (2) *Apply a cravat bandage to the jaw.*

1
2 (a) Place the bandage under the chin and carry its ends upward.
3 Adjust the bandage to make one end longer than the other (Figure 3-20A).

4
5 (b) Take the longer end over the top of the head to meet the short end
6 at the temple and cross the ends over (Figure 3-20B).

7
8 (c) Take the ends in opposite directions to the other side of the head
9 and tie them over the part of the bandage that was applied first (Figure 3-20C).

10
11
12 *Figure 3-20. Applying a cravat bandage to jaw (Illustrated A through C).*

13
14
15 **NOTE**

16 The cravat bandage technique is used to immobilize a fractured
17 jaw or to maintain a sterile dressing that does not have tail bandages
18 attached.

19
20
21 **3-5. Chest Wounds**

22
23 Chest injuries may be caused by accidents, bullet or missile wounds, stab wounds, or falls.
24 These injuries can be serious and may cause death quickly if proper treatment is not given. A
25 casualty with a chest injury may complain of pain in the chest or shoulder area; he may have
26 difficulty with his breathing. His chest may not rise normally when he breathes. The injury may
27 cause the casualty to cough up blood and to have a rapid or a weak heartbeat. A casualty with an
28 open chest wound has a punctured chest wall. The sucking sound heard when he breathes is
29 caused by air leaking into his chest cavity. This particular type of wound is dangerous and will
30 collapse the injured lung (Figure 3-21). Breathing becomes difficult for the casualty because the
31 wound is open. The soldier's life may depend upon how quickly you make the wound airtight.

32
33
34 *Figure 3-21. Collapsed lung.*

35
36
37 **3-6. Chest Wound(s) Procedure**

38
39 *a. Evaluate the Casualty.* Be prepared to perform lifesaving measures. The basic
40 lifesaving measures may include clearing the airway, rescue breathing, chest compressions,
41 treatment for shock, and/or bleeding control.

42
43 *b. Expose the Wound.* If appropriate, cut or remove the casualty's clothing to expose
44 the entire area of the wound. Remember, DO NOT remove clothing that is stuck to the wound
45 because additional injury may result. DO NOT attempt to clean the wound.

1
2 **NOTE**

3 Examine the casualty to see if there is an entry and/or exit wound.
4 If there are two wounds (entry, exit), perform the same procedure
5 for both wounds. Treat the more serious (heavier bleeding, larger)
6 wound first. It may be necessary to improvise a dressing for the
7 second wound by using strips of cloth, such as a torn T-shirt, or
8 whatever material is available. Also, listen for sucking sounds to
9 determine if the chest wall is punctured.

10
11 **CAUTION**

12 If there is an object extending from (impaled in) the wound, DO
13 NOT remove the object. Apply a dressing around the object and
14 use additional improvised bulky materials/dressings (use the cleanest
15 materials available) to build up the area around the object. Apply a
16 supporting bandage over the bulky materials to hold them in place.

17
18 **CAUTION**

19 DO NOT REMOVE protective clothing in a chemical environment.
20 Apply dressings *over* the protective clothing.

21
22 *c. Open the Casualty's Field Dressing Plastic Wrapper.* In cases where there is a
23 sucking chest wound, the plastic wrapper is used with the field dressing to create an airtight seal.
24 If a plastic wrapper is not available, or if an additional wound needs to be treated; cellophane,
25 foil, the casualty's poncho, or similar material may be used. The covering should be wide enough
26 to extend 2 inches or more beyond the edges of the wound in all directions.

27
28 (1) Tear open one end of the casualty's plastic wrapper covering the field
29 dressing. Be careful not to destroy the wrapper and DO NOT touch the inside of the wrapper.

30
31 (2) Remove the inner packet (field dressing).

32
33 (3) Complete tearing open the empty plastic wrapper using as much of the
34 wrapper as possible to create a flat surface.

35
36 *d. Place the Wrapper Over the Wound.* Place the inside surface of the plastic
37 wrapper directly over the wound *when the casualty exhales* and hold it in place (Figure 3-22).
38 The casualty may hold the plastic wrapper in place if he is able.

39
40
41 *Figure 3-22. Open chest wound sealed with plastic wrapper.*

42
43
44 *e. Apply the Dressing to the Wound.*
45

- 1 (1) Use your free hand and shake open the field dressing (Figure 3-23).
2
3

4 *Figure 3-23. Shaking open the field dressing.*
5
6

- 7 (2) Place the white side of the dressing on the plastic wrapper covering the
8 wound (Figure 3-24).
9

10
11 *Figure 3-24. Field dressing placed on plastic wrapper.*
12
13

14 **NOTE**

15 Use the casualty's field dressing, not your own.
16

- 17 (3) Have the casualty breathe normally.
18

- 19 (4) While maintaining pressure on the dressing, grasp one tail of the field
20 dressing with the other hand and wrap it around the casualty's back.
21

- 22 (5) Wrap the other tail in the opposite direction, bringing both tails over the
23 dressing (Figure 3-25).
24
25

26 *Figure 3-25. Tails of field dressing wrapped around casualty in opposite direction.*
27
28

- 29 (6) Tie the tails into a *nonslip* knot in the center of the dressing *after* the
30 casualty exhales and *before* he inhales. This will aid in maintaining pressure on the bandage
31 after it has been tied (Figure 3-26). Tie the dressing firmly enough to secure the dressing without
32 interfering with the casualty's breathing.
33
34

35 *Figure 3-26. Tails of dressing tied into nonslip knot over center of dressing.*
36
37

38 **NOTE**

39 When practical, apply direct manual pressure over the dressing for
40 5 to 10 minutes to help control the bleeding.
41

- 42 *f. Position the Casualty.* Position the casualty on his injured side or in a sitting
43 position, whichever makes breathing easier (Figure 3-27).
44
45

Figure 3-27. Casualty positioned (lying) on injured side.

- g. *Seek Medical Aid.* Contact medical personnel.

WARNING

Even if an airtight dressing has been placed properly, air may still enter the chest cavity without having a means to escape. This causes a life-threatening condition called *tension pneumothorax*. If the casualty's condition (for example, difficulty breathing, shortness of breath, restlessness, or blueness of the skin) worsens after placing the dressing, quickly lift or remove and then replace the airtight dressing.

3-7. Abdominal Wounds

The most serious abdominal wound is one in which an object penetrates the abdominal wall and pierces internal organs or large blood vessels. In these instances, bleeding may be severe and death can occur rapidly.

3-8. Abdominal Wound Procedure

a. *Evaluate the Casualty.* Be prepared to perform basic lifesaving measures. It is necessary to check for both entry and exit wounds. If there are two wounds (entry and exit), treat the wound that appears more serious first (for example, the heavier bleeding, protruding organs, larger wound, and so forth). It may be necessary to improvise dressings for the second wound by using strips of cloth, a T-shirt, or the cleanest material available.

b. *Position the Casualty.* Place and maintain the casualty on his back with his knees in an upright (flexed) position (Figure 3-28). The knees-up position helps relieve pain, assists in the treatment of shock, prevents further exposure of the bowel (intestines) or abdominal organs, and helps relieve abdominal pressure by allowing the abdominal muscles to relax.

Figure 3-28. Casualty positioned (lying) on back with knees (flexed) up.

- c. *Expose the Wound.*

(1) Remove the casualty's loose clothing to expose the wound. However, DO NOT attempt to remove clothing that is stuck to the wound; removing it may cause further injury.

CAUTION

DO NOT REMOVE protective clothing in a chemical environment.
Apply dressings *over* the protective clothing.

(2) Gently pick up any organs that may be on the ground. Do this with a clean, dry dressing or with the cleanest available material. Place the organs on top of the casualty's abdomen (Figure 3-29).

Figure 3-29. Protruding organs placed near wound.

NOTE

- DO NOT probe, clean, or try to remove any foreign object from the abdomen.
- DO NOT touch with bare hands any exposed organs.
- DO NOT push organs back inside the body.

d. Apply the Field Dressing. Use the casualty's field dressing, not your own. If the field dressing is not large enough to cover the entire wound, the plastic wrapper from the dressing may be used to cover the wound first (placing the field dressing on top). Open the plastic wrapper carefully without touching the inner surface. If necessary, other improvised dressings may be made from clothing, blankets, or the cleanest materials available.

WARNING

If there is an object extending from the wound, DO NOT remove it. Place as much of the wrapper over the wound as possible without dislodging or moving the object. DO NOT place the wrapper over the object.

- (1) Grasp the tails in both hands.
- (2) Hold the dressing with the white side down directly over the wound.
- (3) Pull the dressing open and place it directly over the wound (Figure 3-30). If the casualty is able, he may hold the dressing in place.

Figure 3-30. Dressing placed directly over the wound.

(4) Hold the dressing in place with one hand and use the other hand to wrap one of the tails around the body.

1 (5) Wrap the other tail in the opposite direction until the dressing is
2 completely covered. Leave enough of the tail for a knot.

3
4 (6) Loosely tie the tails with a nonslip knot at the casualty's side (Figure 3-31).
5
6

7 *Figure 3-31. Dressing applied and tails tied with a nonslip knot.*
8
9

10 **WARNING**

11 When the dressing is applied, DO NOT put pressure on the wound or
12 exposed internal parts, because pressure could cause further injury
13 (vomiting, ruptured intestines, and so forth). Therefore, tie the dressing
14 ties (tails) loosely at casualty's side, not directly over the dressing.
15

16 (7) Tie the dressing firmly enough to prevent slipping without applying
17 pressure to the wound site (Figure 3-32).
18
19

20 *Figure 3-32. Field dressing covered with improvised material and loosely tied.*
21
22

23 Field dressings can be covered with improvised reinforcement material (cravats, strips of torn T-
24 shirt, or other cloth) for additional support and protection. Tie improvised bandage on the
25 opposite side of the dressing ties firmly enough to prevent slipping but without applying
26 additional pressure to the wound.
27

28 **CAUTION**

29 DO NOT give casualties with abdominal wounds food nor water
30 (moistening the lips is allowed).
31

32 *e. Seek Medical Aid.* Notify medical personnel.
33
34

35 **3-9. Burn Injuries**

36 Burns often cause extreme pain, scarring, or even death. Proper treatment will minimize further
37 injury of the burned area. Before administering the proper first aid, you must be able to
38 recognize the type of burn to be treated. There are four types of burns:
39

40 • Thermal burns caused by fire, hot objects, hot liquids, and gases or by nuclear
41 blast or fireball.
42

43 • Electrical burns caused by electrical wires, current, or lightning.
44

1 • Chemical burns caused by contact with wet or dry chemicals or white phosphorus
2 (WP)--from marking rounds and grenades.

3
4 • Laser burns.
5
6

7 **3-10. First Aid for Burns**

8
9 a. *Eliminate the Source of the Burn.* The source of the burn must be eliminated
10 before any evaluation or treatment of the casualty can occur.
11

12 (1) Remove the casualty quickly from danger and cover the *thermal burn* with
13 any large nonsynthetic material, such as a field jacket. If the casualty's clothing is still on fire,
14 roll the casualty on the ground to smother (put out) the flames (Figure 3-33).
15
16

17 *Figure 3-33. Casualty covered and rolled on ground.*
18
19

20 **CAUTION**

21 Synthetic materials, such as nylon, may melt and cause further injury.
22

23 (2) Remove the *electrical burn* casualty from the electrical source by turning
24 off the electrical current. DO NOT attempt to turn off the electricity if the source is not close by.
25 Speed is critical, so DO NOT waste unnecessary time. If the electricity cannot be turned off,
26 wrap any *nonconductive* material (*dry rope, dry clothing, dry wood, and so forth*) around the
27 casualty's back and shoulders and drag the casualty away from the electrical source (Figure 3-34).
28 DO NOT make body-to-body contact with the casualty or touch any wires because you could
29 also become an *electrical burn casualty*.
30
31

32 *Figure 3-34. Casualty removed from electrical source (using nonconductive material).*
33
34

35 **WARNING**

36 High voltage electrical burns may cause temporary unconsciousness,
37 difficulties in breathing, or difficulties with the heart (heartbeat).
38

39 (3) Remove the *chemical* from the *burned casualty*. Remove *liquid* chemicals
40 by flushing with as much water as possible. Remove *dry* chemicals by brushing off loose
41 particles (DO NOT use the bare surface of your hand because you could become a chemical burn
42 casualty) and then flush with large amounts of water, if available. If *large* amounts of water are
43 not available, then NO water should be applied because small amounts of water applied to a *dry*
44 chemical burn may cause a chemical reaction. When WP strikes the skin, smother with water, a

1 wet cloth, or wet mud. Keep WP covered with a wet material to *exclude* air that will prevent the
2 particles from burning.
3

4 (4) Remove the *laser burn* casualty from the source. (NOTE: Lasers produce
5 a narrow amplified beam of light. The word *laser* means light amplification by stimulated
6 emission of radiation and sources include range finders, weapons/guidance, communication
7 systems, and weapons simulations such as MILES [Multiple Integrated Laser Engagement
8 System].) When removing the casualty from the laser beam source, be careful not to enter the
9 beam or you may become a casualty. Never look directly at the beam source and if possible,
10 wear appropriate eye protection.
11

12 **NOTE**

13 After the casualty is removed from the source of the burn, he should
14 be evaluated for conditions requiring basic lifesaving measures.
15

16 *b. Expose the Burn.* Cut and gently lift away any clothing covering the burned area,
17 without pulling clothing over the burns. Leave in place any clothing that is stuck to the burn. If
18 the casualty's hands or wrists have been burned, remove jewelry if possible without causing
19 further injury (rings, watches, and so forth) and place in his pockets. This prevents the necessity
20 to cut off jewelry since swelling usually occurs as a result of a burn.
21

22 **CAUTION**

- 23 • DO NOT lift or cut away clothing if in a chemical environment.
24 Apply the dressing directly over the casualty's protective clothing.
- 25 • DO NOT attempt to decontaminate skin where blisters have formed.
26

27 *c. Apply a Field Dressing to the Burn.*
28

- 29 (1) Grasp the tails of the casualty's dressing in both hands.
30
- 31 (2) Hold the dressing directly over the wound with the white (sterile) side
32 down, pull the dressing open, and place it directly over the wound. If the casualty is able, he may
33 hold the dressing in place.
34
- 35 (3) Hold the dressing in place with one hand and use the other hand to wrap
36 one of the tails around the limbs or the body.
37
- 38 (4) Wrap the other tail in the opposite direction until the dressing is
39 completely covered.
40
- 41 (5) Tie the tails into a knot over the outer edge of the dressing. The dressing
42 should be applied lightly over the burn. Ensure that dressing is applied firmly enough to prevent
43 it from slipping.
44

NOTE

Use the cleanest improvised dressing material available if a field dressing is not available or if it is not large enough for the entire wound.

d. Take the Following Precautions:

- DO NOT place the dressing over the face or genital area.
- DO NOT break the blisters.
- DO NOT apply grease or ointments to the burns.
- For electrical burns, check for both an entry and exit burn from the passage of electricity through the body. Exit burns may appear on any area of the body despite location of entry burn.
- For burns caused by wet or dry chemicals, flush the burns with large amounts of water and cover with a dry dressing.
- For burns caused by WP, flush the area with water, then cover with a wet material, dressing, or mud to exclude the air and keep the WP particles from burning.
- For laser burns, apply a field dressing.
- If the casualty is conscious and not nauseated, give him small amounts of water.

e. Seek Medical Aid. Notify medical personnel.

3-11. Shoulder Bandage

a. To apply bandages attached to the field first aid dressing—

(1) Take one bandage across the chest and the other across the back and under the arm opposite the injured shoulder.

(2) Tie the ends with a nonslip knot (Figure 3-35).

Figure 3-35. Shoulder bandage.

b. To apply a cravat bandage to the shoulder or armpit—

1
2 (1) Make an extended cravat bandage by using two triangular bandages
3 (Figure 3-36A); place the end of the first triangular bandage along the base of the second one
4 (Figure 3-36B).

5
6 (2) Fold the two bandages into a single extended bandage (Figure 3-36C).

7
8 (3) Fold the extended bandage into a single cravat bandage (Figure 3-36D).
9 After folding, secure the thicker part (overlap) with two or more safety pins (Figure 3-36E).

10
11 (4) Place the middle of the cravat bandage under the armpit so that the front
12 end is longer than the back end and safety pins are on the outside (Figure 3-36F).

13
14 (5) Cross the ends on top of the shoulder (Figure 3-36G).

15
16 (6) Take one end across the back and under the arm on the opposite side and
17 the other end across the chest. Tie the ends (Figure 3-36H).

18
19
20 *Figure 3-36. Extended cravat bandage applied to shoulder (or armpit)*
21 *(Illustrated A through H).*

22
23
24 Be sure to place sufficient wadding in the armpit. DO NOT tie the cravat bandage too tightly.
25 Avoid compressing the major blood vessels in the armpit.

26 27 28 **3-12. Elbow Bandage**

29
30 To apply a cravat bandage to the elbow—

31
32 *a.* Bend the arm at the elbow and place the middle of the cravat at the point of the
33 elbow bringing the ends upward (Figure 3-37A).

34
35 *b.* Bring the ends across, extending both downward (Figure 3-37B).

36
37 *c.* Take both ends around the arm and tie them with a nonslip knot at the front of the
38 elbow (Figure 3-37C).

39
40
41 *Figure 3-37. Elbow bandage (Illustrated A through C).*

CAUTION

If an elbow fracture is suspected, DO NOT bend the elbow; bandage it in the position found.

3-13. Hand Bandage

a. To apply a triangular bandage to the hand—

(1) Place the hand in the middle of the triangular bandage with the wrist at the base of the bandage (Figure 3-38A). Ensure that the fingers are separated with absorbent material to prevent chafing and irritation of the skin.

(2) Place the apex over the fingers and tuck any excess material into the pleats on each side of the hand (Figure 3-38B).

(3) Cross the ends on top of the hand, take them around the wrist, and tie them (Figures 3-38C, D, and E) with a nonslip knot.

Figure 3-38. Triangular bandage applied to hand (Illustrated A through E).

b. To apply a cravat bandage to the palm of the hand—

(1) Lay the middle of the cravat over the palm of the hand with the ends hanging down on each side (Figure 3-39A).

(2) Take the end of the cravat at the little finger across the back of the hand, extending it upward over the base of the thumb; then bring it downward across the palm (Figure 3-39B).

(3) Take the thumb end across the back of the hand, over the palm, and through the hollow between the thumb and palm (Figure 3-39C).

(4) Take the ends to the back of the hand and cross them; then bring them up over the wrist and cross them again (Figure 3-39D).

(5) Bring both ends down and tie them with a nonslip knot on top of the wrist (Figure 3-39E and F).

Figure 3-39. Cravat bandage applied to palm of hand (Illustrated A through F).

1 **3-14. Leg (Upper and Lower) Bandage**
2

3 To apply a cravat bandage to the leg—
4

- 5 a. Place the center of the cravat over the dressing (Figure 3-40A).
6
7 b. Take one end around and up the leg in a spiral motion and the other end around
8 and down the leg in a spiral motion, overlapping part of each preceding turn (Figure 3-40B).
9
10 c. Bring both ends together and tie them (Figure 3-40C) with a nonslip knot.
11
12

13 *Figure 3-40. Cravat bandage applied to leg (Illustrated A through C).*
14
15

16 **3-15. Knee Bandage**
17

18 To apply a cravat bandage to the knee as illustrated in Figure 3-41, use the same technique
19 applied in bandaging the elbow.
20

21 **CAUTION**

22 If a fracture of the kneecap is suspected, DO NOT bend the knee;
23 bandage it in the position found.
24
25

26 *Figure 3-41. Cravat bandage applied to knee (Illustrated A through C).*
27
28

29 **3-16. Foot Bandage**
30

31 To apply a triangular bandage to the foot—
32

- 33 a. Place the foot in the middle of the triangular bandage with the heel well forward
34 of the base (Figure 3-42A). Ensure that the toes are separated by absorbent material to prevent
35 chafing and irritation of the skin.
36
37 b. Place the apex over the top of the foot and tuck any excess material into the pleats
38 on each side of the foot (Figure 3-42B).
39
40 c. Cross the ends on top of the foot, take them around the ankle, and tie them at the
41 front of the ankle (Figure 3-42C, D, and E).
42
43

44 *Figure 3-42. Triangular bandage applied to foot (Illustrated A through E).*

CHAPTER 4

FIRST AID FOR FRACTURES

4-1. General

A fracture is any break in the continuity of a bone. Fractures can cause total disability or in some cases death by severing vital organs and/or arteries. On the other hand, they can most often be treated so there is a complete recovery. The potential for recovery depends greatly upon the first aid the individual receives before he is moved. First aid includes immobilizing the fractured part in addition to applying lifesaving measures when necessary. *The basic splinting principle is to immobilize the joints above and below the fracture.*

4-2. Kinds of Fractures

Figure 4-1 depicts types of fractures.

Figure 4-1. Types of fractures (Illustrated A through C).

a. Closed Fracture. A closed fracture is a broken bone that does not break the overlying skin. The tissue beneath the skin may be damaged. A *dislocation* is when a joint, such as a knee, ankle, or shoulder, is not in the proper position. A *sprain* is when the connecting tissues of the joints have been torn. Dislocations and sprains should be treated as closed fractures.

b. Open Fracture. An open fracture is a broken bone that breaks (pierces) the overlying skin. The broken bone may come through the skin or a missile such as a bullet or shell fragment may go through the flesh and break the bone. An open fracture is contaminated and subject to infection.

4-3. Signs and Symptoms of Fractures

Indications of a fracture are deformity, tenderness, swelling, pain, inability to move the injured part, protruding bone, bleeding, or discolored skin at the injury site. A sharp pain when the soldier attempts to move the part is also a sign of a fracture. DO NOT encourage the casualty to move the injured part in order to identify a fracture since such movement could cause further damage to surrounding tissues and promote shock. If you are not sure whether a bone is fractured, care for the injury as a fracture.

4-4. Purposes of Immobilizing Fractures

A fracture is immobilized to prevent the sharp edges of the bone from moving and cutting tissue, muscle, blood vessels, and nerves. This reduces pain and helps prevent or control shock. In a closed fracture, immobilization keeps bone fragments from causing an open wound which can become contaminated and subject to infection.

4-5. Splints, Padding, Bandages, Slings, and Swathes

a. Splints. Splints may be improvised from such items as boards, poles, sticks, tree limbs, or cardboard. If nothing is available for a splint, the chest wall can be used to immobilize a fractured arm and the uninjured leg can be used to immobilize (to some extent) the fractured leg.

b. Padding. Padding may be improvised from such items as a jacket, blanket, poncho, shelter half, or leafy vegetation.

c. Bandages. Bandages may be improvised from belts, rifle slings, bandoliers, kerchiefs, or strips torn from clothing or blankets. Narrow materials such as wire or cord should not be used to secure a splint in place.

d. Slings. A sling is a bandage suspended from the neck to support an upper extremity. If a bandage is not available, a sling can be improvised by using the tail of a coat or shirt or pieces of cloth torn from such items as clothing and blankets. The triangular bandage is ideal for this purpose. Remember that the casualty's hand should be higher than his elbow, and the sling should be applied so that the supporting pressure is on the uninjured side.

e. Swathes. Swathes are any bands (pieces of cloth or load bearing equipment [LBE]) that are used to further immobilize a splinted fracture. Triangular and cravat bandages are often used and are called *swathe bandages*. The purpose of the swathe is to immobilize; therefore, the swathe bandage is placed above and/or below the fracture---not over it.

4-6. Procedures for Splinting Suspected Fractures

Before beginning first aid procedures for a fracture, gather whatever splinting materials are available. Ensure that splints are long enough to immobilize the joint above and below the suspected fracture. If possible, use at least four ties (two above and two below the fracture) to secure the splints. The ties should be nonslip knots and should be tied away from the body on the splint.

a. Evaluate the Casualty. Be prepared to perform any necessary lifesaving measures. Monitor the casualty for development of conditions which may require you to perform necessary lifesaving measures.

WARNING

Unless there is immediate life-threatening danger, such as a fire or an explosion, **DO NOT** move the casualty with a suspected back or neck injury. Improper movement may cause permanent injury.

WARNING

In a chemical environment, **DO NOT** remove any protective clothing. Apply the dressings and splints over the garments.

b. Locate the site of the suspected fracture.

(1) Ask the casualty for the location of the injury.

- Does he have any pain?
- Where is it tender?
- Can he move the extremity?

(2) Look for an unnatural position of the extremity.

(3) Look for a bone sticking out (protruding).

c. Prepare the Casualty for Splinting the Suspected Fracture.

(1) Reassure the casualty. Tell him that you will be providing first aid for him and that medical help is on the way.

(2) Loosen any tight or binding clothing.

(3) Remove all jewelry from the injured part and place it in the casualty's pocket. Tell the casualty you are doing this because if the jewelry is not removed and swelling occurs later, he may not be able to get it off and further bodily injury could result.

(4) Boots should not be removed from the casualty unless they are needed to stabilize a neck injury or there is actual bleeding from the foot.

d. Gather Splinting Materials. If standard splinting materials (splints, padding, and cravats) are not available, gather improvised materials. If splinting material is not available and the suspected fracture **CAN NOT** be splinted, then swathes, or a combination of swathes and lings can be used to immobilize the extremity.

e. Pad the Splints. Pad the splints where they touch any bony part of the body, such as the elbow, wrist, knee, ankle, crotch, or armpit areas. Padding prevents excessive pressure on the area.

f. Check the Circulation Below the Site of the Injury.

(1) Note any pale, white, or bluish-gray color of the skin, which may indicate impaired circulation. Circulation can also be checked by depressing the toe or fingernail beds and observing how quickly the color returns. A slower return of color to the injured side when compared with the uninjured side indicates a problem with circulation. Depressing the toe or fingernail beds is a method to use to check the circulation in a dark-skinned casualty.

(2) Check the temperature of the injured extremity. Use your hand to compare the temperature of the injured side with the uninjured side. The body area below the injury may be colder to the touch indicating poor circulation.

(3) Question the casualty about the presence of numbness, tightness, cold, or tingling sensations.

WARNING

Casualties with fractures of the extremities may show impaired circulation, such as numbness, tingling, cold or pale to bluish skin tone. These casualties should be evacuated by medical personnel and treated as soon as possible. Prompt medical treatment may prevent possible loss of the limb.

WARNING

If it is an open fracture and the bone is protruding from the skin, **DO NOT ATTEMPT TO PUSH THE BONE BACK UNDER THE SKIN.** Apply a field dressing over the wound to protect the area.

g. Apply the Splint in Place.

(1) Splint the fracture in the position found. **DO NOT** attempt to reposition or straighten the injury. If it is an open fracture, stop the bleeding and protect the wound. Cover all wounds with field dressings before applying a splint. Remember to use the casualty's field dressing, not your own.

(2) Place one splint on each side of the fracture. Make sure that the splints reach, if possible, beyond the joints above and below the fracture.

(3) Tie the splints. Secure each splint in place above and below the fracture site with improvised (or actual) cravats. Improvised cravats, such as strips of cloth, belts, or whatever else you have, may be used. With minimal motion to the injured areas, place and tie the splints with the bandages. Push cravats through and under the natural body curvatures, and then gently position improvised cravats and tie in place. Use nonslip knots. Tie all knots on the splint away from the casualty (Figure 4-2). **DO NOT** tie cravats directly over the suspected fracture site.

Figure 4-2. Nonslip knots tied away from casualty.

h. Check the Splint for Tightness.

(1) **CHECK** to be sure that bandages are tight enough to securely hold splinting materials in place, but not so tight that circulation is impaired.

(2) **RECHECK** the circulation after application of the splint. Check the skin color and temperature. This is to ensure that the bandages holding the splint in place have not been tied too tightly. A fingertip check can be made by inserting the tip of the finger between the wrapped tails and the skin.

(3) **MAKE** any necessary adjustment without allowing the splint to become ineffective.

i. Apply a Sling. An improvised sling may be made from any available nonstretching piece of cloth, such as a battle dress uniform (BDU) shirt or trousers, poncho, or shelter half. Slings may also be improvised using the tail of a coat, belt, or a piece of cloth. Figure 4-3 depicts a shirttail used for support. A trousers belt or LBE may also be used for support (Figure 4-4). A sling should place the supporting pressure on the casualty's uninjured side. The supported arm should have the hand positioned slightly higher than the elbow.

Figure 4-3. Shirttail used for support.

Figure 4-4. Belt used for support.

(1) Insert the splinted arm in the center of the sling (Figure 4-5).

Figure 4-5. Arm inserted in center of improvised sling.

(2) Bring the ends of the sling up and tie them at the side (or hollow) of the neck on the uninjured side (Figure 4-6).

Figure 4-6. Ends of improvised sling tied to side of neck.

- (3) Twist and tuck the corner of the sling at the elbow (Figure 4-7).

Figure 4-7. Corner of sling twisted and tucked at elbow.

j. Apply a Swathe. You may use any large piece of cloth, such as a soldier's belt or pistol belt, to improvise a swathe.

WARNING

The swathe should not be placed directly on top of the injury, but positioned either above or below the fracture site.

- (1) Apply swathes to the injured arm by wrapping the swathe over the injured arm, around the casualty's back and under the arm on the uninjured side. Tie the ends on the uninjured side (Figure 4-8).

Figure 4-8. Arm immobilized with strip of clothing.

- (2) A swathe is applied to an injured leg by wrapping the swathe around both legs and securing it on the uninjured side.

k. Seek Medical Aid. Notify medical personnel, watch closely for development of life-threatening conditions.

4-7. Upper Extremity Fractures

Figures 4-9 through 4-16 show how to apply slings, splints, and cravats (swathes) to immobilize and support fractures of the upper extremities. *Although the padding is not visible in some of the illustrations, it is always preferable to apply padding along the injured part for the length of the splint and especially where it touches any bony parts of the body.*

Figure 4-9. Application of triangular bandage to form sling (two methods).

Figure 4-10. Completing sling sequence by twisting and tucking the corner of the sling at the elbow.

Figure 4-11. Board splints applied to fractured elbow when elbow is not bent (two methods).

Figure 4-12. Chest wall used as splint for upper arm fracture when no splint is available.

Figure 4-13. Chest wall, sling, and cravat used to immobilize fractured elbow when elbow is bent.

Figure 4-14. Board splint applied to fractured forearm.

Figure 4-15. Fractured forearm or wrist splinted with sticks and supported with tail of shirt and strips of material.

Figure 4-16. Board splint applied to fractured wrist and hand.

4-8. Lower Extremity Fractures

Figures 4-17 through 4-22 show how to apply splints to immobilize fractures of the lower extremities. *Although padding is not visible in some of the illustrations, it is preferable to apply padding along the injured part for the length of the splint and especially where it touches any bony parts of the body.*

Figure 4-17. Board splints applied to fractured hip or thigh.

Figure 4-18. Board splint applied to fractured or dislocated knee.

Figure 4-19. Board splints applied to fractured lower leg or ankle.

Figure 4-20. Improvised splints applied to fractured lower leg or ankle.

Figure 4-21. Poles rolled in a blanket and used as splints applied to fractured lower extremity.

Figure 4-22. Uninjured leg used as splint for fractured leg (anatomical splint).

4-9. Jaw, Collarbone, and Shoulder Fractures

a. Apply a cravat to immobilize a fractured jaw as illustrated in Figure 4-23. Direct all bandaging support to the top of the casualty's head, not to the back of his neck. If incorrectly placed, the bandage will pull the casualty's jaw back and interfere with his breathing.

Figure 4-23. Fractured jaw immobilized.

WARNING

Casualties with lower jaw (mandible) fractures cannot be laid flat on their backs because facial muscles will relax and may cause an airway obstruction.

b. Apply two belts, a sling, and a cravat to immobilize a fractured collarbone, as illustrated in Figure 4-24.

Figure 4-24. Application of belts, sling, and cravat to immobilize a fractured collarbone.

c. Apply a sling and a cravat to immobilize a fractured or dislocated shoulder, using the technique illustrated in Figure 4-25.

Figure 4-25. Application of sling and cravat to immobilize a fractured or dislocated shoulder.

4-10. Spinal Column Fractures

It is often impossible to be sure a casualty has a fractured spinal column. Be suspicious of any back injury, especially if the casualty has fallen or if his back has been sharply struck or bent. If a casualty has received such an injury and does not have feeling in his legs or cannot move them, you can be reasonably sure that he has a severe back injury which should be managed as a fracture. Remember, if the spine is fractured, bending it can cause the sharp bone fragments to bruise or cut the spinal cord and result in permanent paralysis (Figure 4-26a). The spinal column must maintain a swayback position to remove pressure from the spinal cord.

- a. If the casualty is not to be transported until medical personnel arrive—
- Caution him not to move. Ask him if he is in pain or if he is unable to move any part of his body.
 - Leave him in the position in which he is found. *DO NOT* move any part of his body, unless he is in imminent danger.
 - Slip a blanket or material of similar size under the arch of his back to support the spinal column in a swayback position (Figure 4-26b), if he is in a face up position. If he is lying face down, *DO NOT* put anything under any part of his body.

Figure 4-26. Spinal column must maintain a swayback position.

- b. If the casualty must be transported to a safe location before medical personnel arrive and if the casualty is in a—
- Face-up position, transport him by litter or use a firm substitute, such as a wide board or a door longer than his height. Loosely tie the casualty's wrists together over his waistline, using a cravat or a strip of cloth. Tie his feet together to prevent the accidental dropping or shifting of his legs. Lay a folded blanket across the litter where the arch of his back is to be placed. Using a four-man team (Figure 4-27), place the casualty on the litter without bending his spinal column or his neck.

Figure 4-27. Placing face-up casualty with fractured back onto litter.

- The numbers two, three, and four men position themselves on one side of the casualty; all kneel on one knee along the side of the casualty. The number one man positions himself to the opposite side of the casualty. The numbers two, three, and four men gently place their hands under the casualty. The number one man on the opposite side places his hands under the injured part to assist.

- When all four men are in position to lift, the number two man commands, "PREPARE TO LIFT" and then, "LIFT." All men, in unison, gently lift the casualty about 8 inches. Once the casualty is lifted, the number one man recovers and slides the litter under the casualty, ensuring that the blanket is in proper position. The number one man then returns to his original lift position (Figure 4-27).

- When the number two man commands, "LOWER CASUALTY," all men, in unison, gently lower the casualty onto the litter.

- Facedown position, he must be transported in this same position. The four-man team lifts him onto a regular or improvised litter, keeping the spinal column in a swayback position. If a regular litter is used, first place a folded blanket on the litter at the point where the chest will be placed.

4-11. Neck Fractures

A fractured neck is extremely dangerous. Bone fragments may bruise or cut the spinal cord just as they might in a fractured back.

- a. If the casualty is not to be transported until medical personnel arrive—
 - Caution him not to move. Moving may cause permanent injury or death.
 - Leave the casualty in the position in which he is found. If his neck and head are in an abnormal position, immediately immobilize the neck and head.
 - Keep his head still, if the casualty is lying face up, raise his shoulders slightly, and slip a roll of cloth that has the bulk of a bath towel under his neck (Figure 4-28). The roll should be thick enough to arch his neck only slightly, leaving the back of his head on the ground. DO NOT bend his neck or head forward. DO NOT raise or twist his head. Immobilize the casualty's head (Figure 4-29). Do this by padding heavy objects (such as rocks or his boots filled with dirt, sand, gravel, or rock) and placing them on each side of his head. If it is necessary to use boots, after filling them, tie the top tightly or stuff with pieces of cloth to secure the contents.)

Figure 4-28. Casualty with roll of cloth (bulk) under neck.

Figure 4-29. Immobilization of fractured neck.

- DO NOT move him if the casualty is lying face down. Immobilize the head and neck by padding heavy objects and placing them on each side of his head. DO NOT put a roll of cloth under the neck. DO NOT bend the neck or head, nor roll the casualty onto his back.

b. If the casualty must be prepared for transportation before medical personnel arrive—

NOTE

If the casualty has a fractured neck, at least two persons are needed because the casualty's head and trunk must be moved in unison. The two persons must work in close coordination (Figure 4-30) to avoid bending the neck.

- A wide board is placed lengthwise beside the casualty. It should extend at least 4 inches beyond the casualty's head and feet (Figure 4-30a).

- If the casualty is lying face up, the number one man steadies the casualty's head and neck between his hands. At the same time, the number two man positions one foot and one knee against the board to prevent it from slipping. He then grasps the casualty underneath his shoulder and hip and gently slides him onto the board (Figure 4-30b).

- If the casualty is lying face down, the number one man steadies the casualty's head and neck between his hands, while the number two man gently rolls the casualty over onto the board (Figure 4-30c).

- The number one man continues to steady the casualty's head and neck. The number two man simultaneously raises the casualty's shoulders slightly, places padding under his neck, and immobilizes the casualty's head (Figures 4-30d and e).

- Any improvised supports are secured in position with a cravat or strip of cloth extended across the casualty's forehead and under the board (Figure 4-30d).

- The board is lifted onto a litter or blanket in order to transport the casualty (Figure 4-30e).

Figure 4-30. Preparing casualty with fractured neck for transportation.

CHAPTER 5

FIRST AID FOR CLIMATIC INJURIES

5-1. General

a. It is desirable, but not always possible, for an individual's body to become adjusted (acclimated) to an environment.

(1) The soldier's physical condition determines the amount of time the body needs to adjust to the environment. Even those individuals in good physical condition need time before working or training in extremes of hot or cold weather. Climate-related injuries are usually preventable; prevention is both an individual and leadership responsibility.

(2) Several factors contribute to health and well-being in any environment--

- Diet.
- Sleep and rest.
- Exercise.
- Suitable clothing.

(3) Diet should be suited to an individual's needs in a particular climate. A special diet undertaken for any purpose (such as weight reduction) should be done so with appropriate medical supervision. Specialized clothing and equipment (such as cold weather gear) for a specific environment should be obtained and used properly.

b. For information on the prevention of heat and cold injuries, refer to Field Manual (FM) 21-10.

5-2. Heat Injuries

a. Heat injuries are environmental injuries. They may result when a soldier—

- Is exposed to extreme heat, such as from the sun or from high temperatures.
- Does not wear proper clothing.
- Is in MOPP gear.

- Is inside closed spaces, such as inside an armored vehicle.
 - Wears body armor.
- b.* Heat injury can be divided into three categories: heat cramps, heat exhaustion, and heatstroke.
- c.* Each soldier must be able to recognize and give first aid for heat injuries.

WARNING

The heat casualty should be continually monitored for development of conditions which may require the performance of necessary basic lifesaving measures.

CAUTION

Do not use salt solutions in first aid procedures for heat injuries.

- (1) Check the casualty for signs and symptoms of *heat cramps*.

- **Signs and symptoms.** Heat cramps are caused by an imbalance of chemicals (called electrolytes) in the body as a result of excessive sweating. This condition causes the casualty to exhibit:

- Muscle cramps in the extremities (arms and legs).
- Abdominal (stomach) cramps.
- Excessive sweating.

NOTE

Thirst may or may not occur. Heat cramps can occur without the soldier being thirsty.

- **First aid measures.**
 - Move the casualty to a cool, shady area or improvise shade if none is available.
 - Loosen his clothing (if not in a chemical environment).

NOTE

In a chemical environment, transport the heat casualty to a noncontaminated area as soon as the mission permits.

- Have him slowly drink at least one canteen full of water.
(The body absorbs cool water faster than warm or cold water; therefore, cool water is preferred if it is available.)

- Seek medical aid should cramps continue.

(2) Check the casualty for signs and symptoms of *heat exhaustion*.

- **Signs and symptoms.** Heat exhaustion is caused by loss of water (dehydration) through sweating without adequate fluid replacement. It can occur in an otherwise fit individual who is involved in physical exertion in any hot environment especially if the soldier is not acclimatized to that environment. These signs and symptoms are—

- Excessive sweating with pale, moist, cool skin.
- Headache.
- Weakness.
- Dizziness.
- Loss of appetite.
- Heat cramps.
- Nausea (with or without vomiting).
- Urge to defecate.
- Chills (gooseflesh).
- Rapid breathing.
- Tingling of hands and/or feet.
- Confusion.

- **First aid measures.**

- Move the casualty to a cool, shady area or improvise shade if none is available.

- Loosen or remove his clothing and boots and pour water on him and fan him (unless in a chemical environment).

- Have him slowly drink at least one canteen of water.
- Elevate his legs.
- If possible, the casualty should not participate in strenuous activity for the remainder of the day.
- Monitor the casualty until the symptoms are gone, or medical aid arrives.

(3) Check the casualty for signs and symptoms of *heatstroke*.

WARNING

Heatstroke is a medical emergency which may result in death if care is delayed.

- **Signs and symptoms.** A soldier suffering from heatstroke has been exposed to high temperatures (such as direct sunlight or being dressed in protective overgarments). Heatstroke occurs more rapidly in soldiers who are engaged in work or other physical activity in a high heat environment. Heatstroke is caused by a failure of the body's cooling mechanism which includes a decrease in the body's ability to produce sweat. The casualty's skin is red (flushed), hot, and dry. He may experience weakness, dizziness, confusion, headaches, seizures, nausea, stomach pains or cramps, and his respiration and pulse may be rapid and weak. Unconsciousness and collapse may occur suddenly.

- **First aid measures.** Cool casualty immediately by—
 - Moving him to a cool, shady area or improvising shade if none is available.
 - Loosening or removing his clothing (except in a chemical environment).
 - Spraying or pouring water on him; fanning him to permit the coolant effect of evaporation.
 - Massaging his extremities and skin which increases the blood flow to those body areas, thus aiding the cooling process.
 - Elevating his legs.
 - Having him slowly drink at least one canteen full of water if he is conscious.

NOTE

Start cooling casualty immediately. Continue cooling while awaiting transportation and during transport to an MTF.

- **Medical aid.** Seek medical aid because the casualty should be transported to an MTF as soon as possible. Do not interrupt the cooling process or lifesaving measures to seek help; if someone else is present send them for the aid. The casualty should be continually monitored for development of conditions which may require the performance of necessary basic lifesaving measures.

d. *Table.* See Table 5-1 for further information.

Table 5-1. Heat Injuries

INJURIES	SIGNS/SYMPTOMS	FIRST AID¹
Heat Cramps	The casualty experiences muscle cramps Of the arms, legs, and/or stomach. The casualty may also have excessive sweating.	<ol style="list-style-type: none"> 1. Move the casualty to a cool shady area or improvise shade and loosen clothing.² 2. Have him slowly drink at least one canteen full of cool water slowly. 3. Monitor the casualty and give him more water as tolerated.
Heat Exhaustion	The casualty experiences heavy sweating with pale, moist, cool skin; headache, weakness, dizziness, and/or loss of appetite. heat cramps, nausea (with or without vomiting), urge to defecate, chills (gooseflesh), rapid breathing, confusion, and tingling of the hands and/or feet.	<ol style="list-style-type: none"> 1. Move the casualty to a cool, shady area or improvise shade and loosen or remove his clothing.² 2. Pour water on him and fan him to permit the coolant effect of evaporation. 3. Have him slowly drink at least one canteen full of cool water.
Heat Stroke³ (Sunstroke)	The casualty stops sweating (red [flushed] hot, dry skin). He first may experience headache, dizziness, nausea, fast pulse and respiration,	<ol style="list-style-type: none"> 1. Move the casualty to a cool, shady area or improvise shade and loosen or remove his clothing, remove the outer garments

	<p>seizures, and mental confusion. He may collapse and suddenly become unconscious. THIS IS A MEDICAL EMERGENCY.</p>	<p>and protective clothing if the situation permits.² 2. Start cooling the casualty immediately. Spray or pour water on him. Fan him. Massage his extremities and skin. 3. Elevate his legs. 4. If conscious, have him slowly drink at least one canteen full of cool water. 5. Seek medical aid. Continue cooling while awaiting transport and continue first aid while en route.</p>
<p>LEGEND: ¹ The first aid procedure for heat related injuries caused by wearing individual protective equipment (IPE) is to move the casualty to a clean area and give him water to drink. ² When in a chemical environment, do not loosen or remove the casualty's clothing. ³ Can be fatal if not provided first aid and medical treatment promptly.</p>		

5-3. Cold Injuries

Cold injuries are most likely to occur when conditions are moderately cold, but accompanied by wet or windy conditions. Cold injuries can usually be prevented. Well-disciplined and well-trained soldiers can be protected even in the most adverse circumstances. They and their leaders must know the hazards of exposure to the cold. They must know the importance of personal hygiene, exercise, care of the feet and hands, and the use of protective clothing.

a. Contributing Factors.

(1) Temperature, humidity, precipitation, and rain greatly increases likelihood of cold injuries, and the soldier with wet clothing are at great risk of cold injuries. Riverine operations (river, swamp, and stream crossings) increase likelihood of cold injuries. Low temperatures and low relative humidity (dry cold) promote frostbite. Higher temperatures, together with moisture, promote immersion syndrome. Windchill accelerates the loss of body heat and may aggravate cold injuries.

(2) Relatively stationary activities such as being in an observation post or on guard duty increase the soldier's vulnerability to cold injury. Also, a soldier is more likely to receive a cold injury if he is—

- In contact with the ground (such as marching, performing guard duty, or engaging in other outside activities).
- Immobile for long periods (such as while riding in an unheated or open vehicle).
- Standing in water, such as in a foxhole.
- Out in the cold for days without being warmed.
- Deprived of an adequate diet and rest.
- Not able to take care of his personal hygiene.

(3) Physical fatigue contributes to apathy, which leads to inactivity, personal neglect, carelessness, and reduced heat production. In turn, these increase the risk of cold injury. Soldiers with prior cold injuries have a higher-than-normal risk of subsequent cold injury, not necessarily involving the body part previously injured.

(4) Depressed or unresponsive soldiers are also vulnerable because they are less active. These soldiers tend to be careless about precautionary measures, especially warming activities, when cold injury is a threat.

(5) Excessive use of alcohol or drugs leading to faulty judgement or unconsciousness in a cold environment increases the risk of becoming a cold injury casualty.

b. Signs and Symptoms. Once a soldier becomes familiar with the factors that contribute to cold injury, he must learn to recognize cold injury signs and symptoms.

(1) Many soldiers suffer cold injury without realizing what is happening to them. They may be cold and generally uncomfortable. These soldiers often do not notice the injured part because it is already numb from the cold.

(2) Superficial cold injury usually can be detected by numbness or tingling sensations. These signs and symptoms often can be relieved simply by loosening boots or other clothing and by exercising to improve circulation. In more advanced cases involving deep cold injury, the soldier often is not aware that there is a problem until the affected part feels like a stump or block of wood.

(3) Outward signs of cold injury include discoloration of the skin at the site of injury. In light-skinned persons, the skin first reddens and then becomes pale or waxy white. In

dark-skinned persons, grayness in the skin is usually evident. An injured foot or hand feels cold to the touch. Swelling may be an indication of deep injury. Also note that blisters may occur after rewarming the affected parts. Soldiers should work in pairs (buddy teams) to check each other for signs of discoloration and other symptoms.

c. First aid measures. First aid for cold injuries depends on whether they are superficial or deep. Cases of superficial cold injury can be adequately treated by rewarming the affected part using body heat. (For example, this can be done by covering cheeks with hands, putting fingertips in armpits, or placing the casualty's feet under the clothing of a buddy [next to his belly].) The injured part should NOT be massaged, exposed to a fire or stove, rubbed with snow, slapped, chafed, or soaked in cold water. Walking on injured feet should be avoided. Deep cold injury (frostbite) is very serious and requires prompt first aid to avoid or to minimize the loss of parts or all of the fingers, toes, hands, or feet. The sequence for treating cold injuries depends on whether the condition is life threatening. The first priority in managing cold injuries is to remove the casualty from the cold environment (such as building an improvised shelter). Other injuries the casualty may have are provided first aid simultaneously while waiting for transportation or evacuation. If the casualty is to be transported in a nonmedical vehicle first aid measures should be continued en route to the MTF.

d. Conditions Caused by Cold. Conditions caused by cold include chilblain, immersion syndrome (immersion foot and trench foot), frostbite, snow blindness, dehydration, and hypothermia.

(1) **Chilblain.**

- *Signs and symptoms.* Chilblain is caused by repeated prolonged exposure of bare skin at temperatures from 60° Fahrenheit (F) to 32°F, or 20°F for acclimated, dry, unwashed skin. The area may be acutely swollen, red, tender, and hot with itchy skin. There may be no loss of skin tissue in untreated cases but continued exposure may lead to infected, ulcerated, or bleeding lesions.

- *First aid measures.* Within minutes, the area usually responds to locally applied body heat. Rewarm the affected part by applying firm steady pressure with your hands, or placing the affected part under your arms or against the stomach of a buddy. DO NOT rub or massage affected areas.

NOTE

Medical personnel should evaluate the injury, because signs and symptoms of tissue damage may be slow to appear.

(2) **Immersion syndrome (immersion foot and trench foot).** Immersion foot and trench foot are injuries that result from fairly long exposure of the feet to wet conditions at temperatures from approximately 32°F to 50°F. Inactive feet in damp or wet socks and boots, or tightly laced boots which impair circulation are even more susceptible to injury. This injury can be very serious; it can lead to loss of toes or parts of the feet. If exposure of the feet has been

prolonged and severe, the feet may swell so much that pressure closes the blood vessels and cuts off circulation. Should an immersion injury occur, dry the feet thoroughly and transport the casualty to an MTF by the fastest means possible.

- *Signs and symptoms.* At first, the parts of the affected foot are cold and painless, the pulse is weak, and numbness may be present. Second, the parts may feel hot, and burning and shooting pains may begin. In later stages, the skin is pale with a bluish cast and the pulse decreases. Other signs and symptoms that may follow are blistering, swelling, redness, heat, hemorrhaging (bleeding), and gangrene.

- *First aid measures.* First aid measures are required for all stages of immersion syndrome injury. Rewarm the injured part gradually by exposing it to warm air. Protect it from trauma and secondary infections. Dry, loose clothing or several layers of warm coverings are preferable to extreme heat. Under no circumstances should the injured part be exposed to an open fire. Elevate the injured part to relieve the swelling. Transport the casualty to an MTF as soon as possible. When the part is rewarmed, the casualty often feels a burning sensation and pain. Symptoms may persist for days or weeks even after rewarming.

NOTE

When providing first aid for immersion foot and trench foot—

- DO NOT massage the injured part.
- DO NOT moisten the skin.
- DO NOT apply heat or ice.

(3) **Frostbite.** Frostbite is the injury of tissue caused from exposure to cold, usually below 32°F depending on the windchill factor, duration of exposure, and adequacy of protection. Individuals with a history of cold injury are likely to suffer an additional cold injury. The body parts most easily frostbitten are the cheeks, nose, ears, chin, forehead, wrists, hands, and feet. Frostbite may involve only the skin (superficial), or it may extend to a depth below the skin (deep). Deep frostbite is very serious and requires prompt first aid to avoid or to minimize the loss of parts or all of the fingers, toes, hands, or feet.

- *Signs and symptoms.*
 - Loss of sensation (numb feeling) in any part of the body.
 - Sudden blanching (whitening) of the skin of the affected part, followed by a momentary tingling sensation.
 - Redness of skin in light-skinned soldiers; grayish coloring in dark-skinned individuals.
 - Blisters.
 - Swelling or tender areas.

- Loss of previous sensation of pain in affected area.
- Pale, yellowish, waxy-looking skin.
- Frozen tissue that feels solid (or wooden) to the touch.

CAUTION

Deep frostbite is a very serious injury and requires immediate first aid and subsequent medical treatment to avoid or minimize loss of body parts.

- *First aid measures.*
 - **Face, ears, and nose.** Cover the casualty's affected area with his and/or your bare hands until sensation and color return.
 - **Hands.** Open the casualty's field jacket and shirt. (In a chemical environment, do not loosen or remove the clothing and protective overgarments.) Place the affected hands under the casualty's armpits. Close the field jacket and shirt to prevent additional exposure.
 - **Feet.** Remove the casualty's boots and socks if he does not need to walk any further to receive additional treatment. (Thawing the casualty's feet and forcing him to walk on them will cause additional pain and injury.) Place the affected feet under clothing and against the body of another soldier.

WARNING

DO NOT attempt to thaw the casualty's feet or other frozen areas if he will be required to walk or travel to receive medical treatment. The casualty should avoid walking, if possible, because there is less danger in walking while the feet are frozen than after they have been thawed. Thawing in the field increases the possibilities of infection, gangrene, or other injury.

NOTE

Thawing may occur spontaneously during transportation to the MTF; this cannot be avoided since the body in general must be kept warm.

In all of the above areas, ensure that the casualty is kept warm and that he is covered (to avoid further injury). Seek medical treatment as soon as possible. Reassure the casualty, protect the affected area from further injury by covering it lightly with a blanket or any dry clothing, and seek shelter out of the wind. Remove or loosen constricting clothing (except in a contaminated environment) and increase insulation. Exercise as much as possible, avoiding trauma to the

injured part, and be prepared for pain when thawing occurs. Protect the frostbitten part from additional injury. DO NOT—

- Rub the injured part with snow or apply cold water soaks.
- Warm the part by massage or exposure to open fire because the frozen part may be burned due to the lack of feeling.
- Use ointments or other salves.
- Manipulate the part in any way to increase circulation.
- Use alcohol or tobacco because this reduces the body's resistance to cold.

NOTE

Remember, when freezing extends to a depth below the skin, it is a much more serious injury. Extra care is required to reduce or avoid the chances of losing all or part of the toes or feet. This also applies to the fingers and hands.

(4) **Snow blindness.** Snow blindness is the effect that glare from an ice field or snowfield has on the eyes. It is more likely to occur in hazy, cloudy weather than when the sun is shining. Glare from the sun will cause an individual to instinctively protect his eyes. However, in cloudy weather, he may be overconfident and expose his eyes longer than when the threat is more obvious. He may also neglect precautions such as the use of protective eyewear. Waiting until discomfort (pain) is felt before using protective eyewear is dangerous because a deep burn of the eyes may already have occurred.

- *Signs and symptoms.* Symptoms of snow blindness are a sensation of grit in the eyes with pain in and over the eyes, made worse by movement the eyeball. Other signs and symptoms are watering, redness, headache, and increased pain on exposure to light.
- *First aid measures.* First aid measures consist of blindfolding or covering the eyes with a dark cloth which stops painful eye movement. Complete rest is desirable. If further exposure to light is not preventable, the eyes should be protected with dark bandages or the darkest glasses available. Once unprotected exposure to sunlight stops, the condition usually heals in a few days without permanent damage. The casualty should be evacuated to the nearest MTF.

(5) **Dehydration.** Dehydration occurs when the body loses too much fluid. A certain amount of body fluid is lost through normal body processes. A normal daily intake of liquids replaces these losses. When individuals are engaged in any strenuous exercises or activities, fluid is lost through sweating and this loss creates an imbalance of fluids in the body, and if not matched by rehydration it can contribute to dehydration. The danger of dehydration is

as prevalent in cold regions as it is in hot regions. In hot weather, the individual is aware of his body losing fluids through sweat. In cold weather, however, it is extremely difficult to realize that this condition exists as sweating is not as apparent as in a hot environment. The danger of dehydration in cold weather operations is a serious problem. In cold climates, sweat evaporates so rapidly or is absorbed so thoroughly by layers of heavy clothing that it is rarely visible on the skin. Dehydration also occurs during cold weather operations because drinking is inconvenient. Dehydration will weaken or incapacitate a casualty for a few hours, or sometimes, several days. Because rest is an important part of the recovery process, casualties must take care that limited movement during their recuperative period does not enhance the risk of becoming a cold injury casualty.

- *Signs and symptoms.* The symptoms of cold weather dehydration are similar to those encountered in heat exhaustion. The mouth, tongue, and throat become parched and dry, and swallowing becomes difficult. The casualty may have nausea (with or without) vomiting along with extreme dizziness and fainting. The casualty may also feel generally tired and weak and may experience muscle cramps. Focusing the eyes may also become difficult.

- *First aid measures.* The casualty should be kept warm and his clothes should be loosened (not in a chemical environment) to allow proper circulation. Shelter from wind and cold must be provided. Fluid replacement should begin immediately and the soldier transported to an MTF as soon as possible.

(6) **Hypothermia (general cooling).** When exposed to prolonged cold weather a soldier may become both mentally and physically numb, thus neglecting essential tasks or requiring more time and effort to achieve them. Under some conditions (particularly cold water immersion), even a soldier in excellent physical condition may die in a matter of minutes. The destructive influence of cold on the body is called *hypothermia*. This means bodies lose heat faster than they can produce it. Hypothermia can occur from exposure to temperatures either above or below freezing, especially from immersion in cold water, wet-cold conditions, or from the effect of wind. Physical exhaustion and insufficient food intake may also increase the risk of hypothermia. General cooling of the entire body to a temperature below 95°F is caused by continued exposure to low or rapidly dropping temperatures, cold moisture, snow, or ice. Fatigue, poor physical condition, dehydration, faulty blood circulation, alcohol or other drug use, trauma, and immersion can cause hypothermia. Remember, cold may affect the body systems slowly and almost without notice. Soldiers exposed to low temperatures for extended periods may suffer ill effects even if they are well protected by clothing.

- *Signs and symptoms.* As the body cools, there are several stages of progressive discomfort and impairment. A sign that is noticed immediately is shivering. Shivering is an attempt by the body to generate heat. The pulse is faint or very difficult to detect. People with temperatures around 90°F may be drowsy and mentally slow. Their ability to move may be hampered, stiff, and uncoordinated, but they may be able to function minimally. Their speech may be slurred. As the body temperature drops further, shock becomes evident as the person's eyes assume a glassy state, breathing becomes slow and shallow, and the pulse becomes

weaker or absent. The person becomes very stiff and uncoordinated. Unconsciousness may follow quickly. As the body temperature drops even lower, the extremities freeze, and a deep (or core) body temperature (below 85°F) increases the risk of irregular heart action. This irregular heart action or heart standstill can result in sudden death.

- *First aid measures.* Except in cases of the most severe hypothermia (marked by coma or unconsciousness and a weak pulse), the first aid measures for hypothermia are directed towards protecting the casualty from further loss of body heat. For the casualty who is conscious, first aid measures are directed at rewarming the body evenly and without delay. Provide heat by using a hot water bottle or field expedient or another soldier's body heat.

CAUTION

DO NOT expose the casualty to an open fire, as he may become burned.

NOTE

When using a hot water bottle or field expedient (canteen filled with warm water), the bottle or canteen must be wrapped in cloth prior to placing it next to the casualty. This will reduce the chance of burning the casualty's skin.

Always call or send for help as soon as possible and protect the casualty immediately with dry clothing or a sleeping bag. Then, move him to a warm place. Evaluate other injuries and provide first aid as required. First aid measures can be performed while the casualty is waiting transportation or while he is en route. In the case of an accidental breakthrough into ice water, or other hypothermic accident, strip the casualty of wet clothing immediately and bundle him into a sleeping bag. Rescue breathing should be started at once if the casualty's breathing has stopped or is irregular or shallow. Warm liquids (**NOT HOT**) may be given gradually if the casualty is conscious. DO NOT force liquids on an unconscious or semiconscious casualty because he may choke. The casualty should be transported on a litter because the exertion of walking may aggravate circulation problems. Medical personnel should immediately treat any hypothermia casualty. Hypothermia is life threatening until normal body temperature has been restored. The first aid measures for a casualty with severe hypothermia are based upon the following principles: attempt to avoid further heat loss, handle the casualty gently, and transport the casualty as soon as possible to the nearest MTF. If at all possible, the casualty should be evacuated by medical personnel.

WARNING

Rewarming a severely hypothermic casualty is extremely dangerous in the field due to the possibility of such complications as rewarming shock and disturbances in the rhythm of the heartbeat. These conditions require treatment by medical personnel.

NOTE

Resuscitation of casualties with hypothermic complications is difficult if not impossible to do outside of an MTF setting.

CAUTION

The casualty is unable to generate his own body heat. Therefore, merely placing him in a blanket or sleeping bag is not sufficient.

e. *Table.* See Table 5-2 for further information.

Table 5-2. Injuries Caused by Cold and Wet Conditions

INJURIES	SIGNS/SYMPTOMS	FIRST AID
Chilblain	Red swollen, hot, tender, itching skin. Continued exposure may lead to infected (ulcerated or bleeding) skin lesions.	<ol style="list-style-type: none"> 1. Area usually response to locally applied rewarming (body heat). 2. Do not rub or message area. 3. Seek medical aid.
Immersion Syndrome (Immersion Foot/Trench Foot)	Affected parts are cold, numb, and painless. Parts may then be hot, with burning and shooting pains. Advanced stage: skin pale with bluish cast; pulse decreases; blistering, swelling, heat, hemorrhaging, and gangrene may follow.	<ol style="list-style-type: none"> 1. Gradual rewarming by exposure to warm air. 2. Do not massage or moisten skin. 3. Protect affected parts from trauma. 4. Dry feet thoroughly, avoid walking. 5. Seek medical aid.
Frostbite	Loss of sensation (numb feeling) in any part of the body. Sudden blanching (whitening) of the skin of the affected part, followed by a momentary tingling sensation. Redness of skin in light-skinned soldiers; grayish coloring in dark-skinned soldiers. Blisters. Swelling or tender areas. Loss of previous sensation of pain in the affected area. Pale yellowish, waxy-	<ol style="list-style-type: none"> 1. Warm the area at the first sign of frostbite, using firm, steady pressure of the hand, underarm, or abdomen. 2. Face, ears, nose: cover area with hands (casualty's own or buddy's). 3. Hands: open field jacket and place casualty's hands against his body, then close the jacket to prevent heat loss. 4. Feet: Remove the

	looking skin. Frozen tissue that feels solid (wooden) to the touch.	casualty's boots and socks and place against the body of another soldier. 5. WARNING: Do not attempt to thaw the casualty's feet or other frozen areas if he will be required to walk or travel to an MTF for additional treatment. The possibility of injury from walking is less when the feet are frozen than when have been thawed. (However, if possible avoid walking.) Thawing in the field increases the possibility of infection, gangrene, or injury. 6. Loosen or remove constricting clothing and remove any jewelry. 7. Increase insulation (cover with blanket or other dry material). Ensure casualty exercises as much as possible, avoiding trauma to injured part.
Snow Blindness	Eyes may feel scratchy. Watery, redness, headache, and increased pain with exposure to light can occur.	1. Cover the eyes with a dark cloth. 2. Seek medical aid.
Dehydration	Similar to heat exhaustion (refer to Table 5-1).	1. Keep warm. 2. Casualty needs fluid replacement, rest, and prompt medical aid.
Hypothermia	Casualty is cold. Shivering. Core temperature is low. Consciousness may be altered. Uncoordinated movements may occur. Shock and coma may result	Mild Hypothermia 1. Rewarm body evenly and without delay. (Need to provide heat source; casualty's body unable to generate heat.)

	as body temperature drops.	<ol style="list-style-type: none">2. Keep dry, protect from the elements.3. Warm (not hot) liquids may be given gradually (to conscious casualties only).4. Be prepared to start basic life support measures for the casualty.5. Seek medical treatment immediately. <p>Severe Hypothermia</p> <ol style="list-style-type: none">1. Stabilize the temperature.2. Attempt to avoid further heat loss.3. Handle the casualty gently.4. Evacuate to the nearest MTF as soon as possible. <p>WARNING: Hypothermia is a medical emergency. Prompt medical attention is necessary.</p>
--	----------------------------	--

CHAPTER 6

FIRST AID FOR BITES AND STINGS

6-1. General

Snakebites, insect bites, or stings can cause intense pain and/or swelling. If not treated promptly and correctly, they can cause serious illness or death. The severity of a snakebite depends upon: whether the snake is poisonous or nonpoisonous, the type of snake, the location of the bite, and the amount of venom injected. Bites from humans and other animals, such as dogs, cats, bats, raccoons, and rats, can cause severe bruises and infection and tears or lacerations of tissue. Awareness of the potential sources of injuries can reduce or prevent them from occurring. Knowledge and prompt application of first aid measures can lessen the severity of injuries from bites and stings and keep the soldier from becoming a serious casualty.

6-2. Types of Snakes

a. Nonpoisonous Snakes. There are approximately 130 different varieties of nonpoisonous snakes in the United States. They have oval-shaped heads and round eyes. Unlike poisonous snakes, discussed below, nonpoisonous snakes do not have fangs with which to inject venom. Figure 6-1 depicts the characteristics of a nonpoisonous snake.

Figure 6-1. Characteristics of nonpoisonous snake.

b. Poisonous Snakes. Poisonous snakes are found throughout the world, primarily in tropical to moderate climates. Within the United States, there are four kinds: rattlesnakes, copperheads, water moccasins (cottonmouth), and coral snakes. Poisonous snakes in other parts of the world include sea snakes, the fer-de-lance, the bushmaster, and the tropical rattlesnake in tropical Central America; the Malayan pit viper in the tropical Far East; the cobra in Africa and Asia; the mamba (or black mamba) in central and southern Africa; and the krait in India and Southeast Asia. Refer to Figure 6-2 for characteristics of a poisonous pit viper.

Figure 6-2. Characteristics of poisonous pit viper.

c. Pit Vipers (Poisonous). Figure 6-3 depicts a variety of poisonous snakes.

Figure 6-3. Poisonous snakes.

(1) Rattlesnakes, bushmasters, copperheads, fer-de-lance, Malayan pit vipers, and water moccasins (cottonmouth) are called pit vipers because of the small, deep pits between the nostrils and eyes on each side of the head (Figure 6-2). In addition to their long, hollow fangs, these snakes have other identifying features: thick bodies, slit-like pupils of the eyes, and flat, almost triangular-shaped heads. Color markings and other identifying characteristics, such as rattles or a noticeable white interior of the mouth (cottonmouth), also help distinguish these poisonous snakes. Further identification is provided by examining the bite pattern of the wound for signs of fang entry. Occasionally there will be only one fang mark, as in the case of a bite on a finger or toe where there is no room for both fangs, or when the snake has broken off a fang.

(2) The casualty's condition provides the best information about the seriousness of the situation, or how much time has passed since the bite occurred. Pit viper bites are characterized by severe burning pain. Discoloration and swelling around the fang marks usually begins within 5 to 10 minutes after the bite. If only minimal swelling occurs within 30 minutes, the bite will almost certainly have been from a nonpoisonous snake or possibly from a poisonous snake which did not inject venom. The venom destroys blood cells, causing a general discoloration of the skin. This reaction is followed by blisters and numbness in the affected area. Other signs which can occur are weakness, rapid pulse, nausea, shortness of breath, vomiting, and shock.

d. Corals, Cobras, Kraits, and Mambas. Corals (Figure 6-4), cobras (Figure 6-5), kraits, and mambas all belong to the same group even though they are found in different parts of the world. All four inject their venom through short, grooved fangs, leaving a characteristic bite pattern.

Figure 6-4. Coral snake.

(1) The small coral snake, found in the Southeastern United States, is brightly colored with bands of red, yellow (or almost white), and black completely encircling the body. Other nonpoisonous snakes have the same coloring, but on the coral snake found in the United States, the red ring always touches the yellow ring. To know the difference between a harmless snake and the coral snake found in the United States, remember the following:

"Red on yellow will kill a fellow,
Red on black, venom will lack."

Figure 6-5. Cobra snake.

(2) The venom of corals, cobras, kraits, and mambas produces symptoms different from those of pit vipers. Because there is only minimal pain and swelling, many people believe that the bite is not serious. Delayed reactions in the nervous system normally occur

between 1 to 7 hours after the bite. Symptoms include blurred vision, drooping eyelids, slurred speech, drowsiness, and increased salivation and sweating. Nausea, vomiting, shock, respiratory difficulty, paralysis, convulsions, and coma will usually develop if the bite is not treated promptly.

e. Sea Snakes. Sea snakes (Figure 6-6) are found in the warm water areas of the Pacific and Indian oceans, along the coasts and at the mouths of some larger rivers. Their venom is VERY poisonous, but their fangs are only 1/4 inch long. The first aid outlined for land snakes also applies to sea snakes.

Figure 6-6. Sea snake.

6-3. Snakebites

a. Poisonous snakes DO NOT always inject venom when they bite or strike a person. However, all snakes may carry tetanus (lockjaw); anyone bitten by a snake, whether poisonous or nonpoisonous, should immediately seek medical attention.

- Poison is injected from the venom sacs through grooved or hollow fangs. Depending on the species, these fangs are either long or short. Pit vipers have long hollow fangs. These fangs are folded against the roof of the mouth and extend when the snake strikes. This allows them to strike quickly and then withdraw. Cobras, coral snakes, kraits, mambas, and sea snakes have short, grooved fangs. These snakes are less effective in their attempts to bite, since they must chew after striking to inject enough venom (poison) to be effective. Figure 6-7 depicts the characteristics of a poisonous snakebite.

- In the event you are bitten, attempt to identify and/or kill the snake. Take it to medical personnel for inspection/identification. This provides valuable information to medical personnel who deal with snakebites. **TREAT ALL SNAKEBITES AS POISONOUS.**

Figure 6-7. Characteristics of poisonous snakebite.

b. The venoms of different snakes cause different effects. Pit viper venom (hemotoxin) destroys tissue and blood cells. Cobras, adders, and coral snakes inject powerful venom (neurotoxin) which affect the central nervous system, causing respiratory paralysis. Water moccasins and sea snakes have venom that is both hemotoxic and neurotoxic.

c. The identification of poisonous snakes is very important since medical treatment will be different for each type of venom. *Unless it can be positively identified, the snake should be killed and saved.* When this is not possible or when doing so is a serious threat to others, identification may sometimes be difficult since many venomous snakes resemble harmless

varieties. When dealing with snakebite problems in foreign countries, seek advice, professional or otherwise, which may help identify species in the particular area of operations.

d. Get the casualty to a medical treatment facility as soon as possible and with minimum movement. Until evacuation or treatment is possible, have the casualty lie quietly and not move any more than necessary. If the casualty has been bitten on an extremity, **DO NOT** elevate the limb; keep the extremity level with the body. Keep the casualty comfortable and reassure him. If the casualty is alone when bitten, he should go to the medical facility himself rather than wait for someone to find him. Unless the snake has been positively identified, attempt to kill it and send it with the casualty. Be sure that retrieving the snake does not endanger anyone or delay transporting the casualty.

(1) If the bite is on an arm or leg, place a constricting band (narrow cravat [swathe], or narrow gauze bandage) one to two fingerbreadths above and below the bite (Figure 6-8). If the bite is on the hand or foot, place a single band above the wrist or ankle. The band should be tight enough to stop the flow of blood near the skin, but not tight enough to interfere with circulation. In other words, it should not have a tourniquet-like affect. If no swelling is seen, place the bands about 1 inch from either side of the bite. If swelling is present, put the bands on the unswollen part at the edge of the swelling. If the swelling extends beyond the band, move the band to the new edge of the swelling. (If possible, leave the old band on, place a new one at the new edge of the swelling, and then remove and save the old one in case the process has to be repeated.)

Figure 6-8. Constricting band.

CAUTION

DO NOT attempt to cut open the bite nor suck out the venom. If the venom should seep through any damaged or lacerated tissues in your mouth, you could immediately lose consciousness or even die.

(2) If the bite is located on an arm or leg, immobilize it at a level below the heart. **DO NOT** elevate an arm or leg even with or above the level of the heart.

CAUTION

When a splint is used to immobilize the arm or leg, take **EXTREME** care to ensure the splinting is done properly and does not bind. Watch it closely and adjust it if any changes in swelling occur.

(3) When possible, clean the area of the bite with soap and water. **DO NOT** use ointments of any kind.

(4) **NEVER** give the casualty food, alcohol, stimulants (coffee or tea), drugs, or tobacco.

- (5) Remove rings, watches, or other jewelry from the affected limb.

6-4. Human and Other Animal Bites

Human or other land animal bites may cause lacerations or bruises. In addition to damaging tissue, bites always present the possibility of infection.

a. Human Bites. Human bites that break the skin may become seriously infected since the mouth is heavily contaminated with bacteria. All human bites **MUST** be treated by medical personnel.

b. Animal Bites. Land animal bites can result in both infection and disease. Tetanus, rabies, and various types of fevers can follow an untreated animal bite. Because of these possible complications, the animal causing the bite should, if possible, be captured or killed (without damaging its head) so that it can be tested for disease.

c. First Aid.

- (1) Cleanse the wound thoroughly with soap.
- (2) Flush it well with water.
- (3) Cover it with a sterile dressing.
- (4) Immobilize the injured arm or leg, if appropriate.
- (5) Transport the casualty immediately to an MTF.

NOTE

If unable to capture or kill the animal, provide medical personnel with any information that will help identify it.

6-5. Marine (Sea) Animals

With the exception of sharks and barracuda, most marine animals will not deliberately attack. The most frequent injuries from marine animals are wounds by biting, stinging, or puncturing. Wounds inflicted by marine animals can be very painful, but are rarely fatal.

a. Sharks, Barracuda, and Alligators. Wounds from these marine animals can involve major trauma as a result of bites and lacerations. Bites from large marine animals are potentially the most life threatening of all injuries from marine animals. Major wounds from

these animals can be treated by controlling the bleeding, preventing shock, giving basic life support, splinting the injury, and by securing prompt medical aid.

b. Turtles, Moray Eels, and Corals. These animals normally inflict minor wounds. Treat by cleansing the wound(s) thoroughly and by splinting if necessary.

c. Jellyfish, Portuguese Man-of-War, Anemones, and Others. This group of marine animals inflict injury by means of stinging cells in their tentacles. Contact with the tentacles produces burning pain with a rash and small hemorrhages on the skin. Shock, muscular cramping, nausea, vomiting, and respiratory distress may also occur. Gently remove the clinging tentacles with a towel and wash or treat the area. Use diluted ammonia or alcohol, meat tenderizer, and talcum powder. If symptoms become severe or persist, seek medical aid.

d. Spiny Fish, Urchins, Stingrays, and Cone Shells. These animals inject their venom by puncturing the skin with their spines. General signs and symptoms include swelling, nausea, vomiting, generalized cramps, diarrhea, muscular paralysis, and shock. Deaths are rare. Treatment consists of soaking the wounds in hot water (when available) for 30 to 60 minutes. This inactivates the heat sensitive toxin. In addition, further first aid measures (controlling bleeding, applying a dressing, and so forth) should be carried out as necessary.

CAUTION

Be careful not to scald the casualty with water that is too hot because the pain of the wound will mask the normal reaction to heat.

6-6. Insect (Arthropod) Bites/Stings

An insect bite or sting can cause great pain, allergic reaction, inflammation, and infection. If not treated correctly, some bites/stings may cause serious illness or even death. When an allergic reaction is not involved, first aid is a simple process. In any case, medical personnel should examine the casualty at the earliest possible time. It is important to properly identify the spider, bee, or creature that caused the bite/sting, especially in cases of allergic reaction.

a. Types of Insects. The insects found throughout the world that can produce a bite or sting are too numerous to mention in detail. Commonly encountered stinging or biting insects include brown recluse spiders (Figure 6-9), black widow spiders (Figure 6-10), tarantulas (Figure 6-11), scorpions (Figure 6-12), urticating caterpillars, bees, wasps, centipedes, conenose beetles (kissing bugs), ants, and wheel bugs. Upon being reassigned, especially to overseas areas, take the time to become acquainted with the types of insects to avoid.

Figure 6-9. Brown recluse spider.

Figure 6-10. Black widow spider.

Figure 6-11. Tarantula.

Figure 6-12. Scorpion.

b. Signs and Symptoms. Discussed in paragraphs (1) and (2) below are the most common effects of insect bites/stings. They can occur alone or in combination with the others.

(1) *Less serious.* Commonly seen signs/symptoms are pain, irritation, swelling, heat, redness, and itching. Hives or wheals (raised areas of the skin that itch) may occur. These are the least severe of the allergic reactions that commonly occur from insect bites/stings. They are usually dangerous only if they affect the air passages (mouth, throat, nose, and so forth), which could interfere with breathing. The bites/stings of bees, wasps, ants, mosquitoes, fleas, and ticks are usually not serious and normally produce mild and localized symptoms. A tarantula's bite is usually no worse than that of a bee sting. Scorpions are rare and their stings (except for a specific species found only in the Southwest desert) are painful but usually not dangerous.

(2) *Serious.* Emergency allergic or hypersensitive reactions sometimes result from the stings of bees, wasps, and ants. Many people are allergic to the venom of these particular insects. Bites or stings from these insects may produce more serious reactions, to include generalized itching and hives, weakness, anxiety, headache, breathing difficulties, nausea, vomiting, and diarrhea. Very serious allergic reactions (called *anaphylactic shock*) can lead to complete collapse, shock, and even death. Spider bites (particularly from the black widow and brown recluse spiders) can also be serious. Venom from the black widow spider affects the nervous system. This venom can cause muscle cramps, a rigid, nontender abdomen, breathing difficulties, sweating, nausea, and vomiting. The brown recluse spider generally produces local rather than system-wide problems; however, local tissue damage around the bite can be severe and can lead to an ulcer and even gangrene.

c. First Aid. There are certain principles that apply regardless of what caused the bite/sting. Some of these are—

- If there is a stinger present (for example, from a bee), remove the stinger by scraping the skin's surface with a fingernail or knife. DO NOT squeeze the sac attached to the stinger because it may inject more venom.
- Wash the area of the bite/sting with soap and water (alcohol or an antiseptic may also be used) to help reduce the chances of an infection and remove traces of venom.
- Remove jewelry from bitten extremities because swelling may occur.

- In most cases of insect bites the reaction will be mild and localized; use ice or cold compresses (if available) on the site of the bite/sting. This will help reduce swelling, ease the pain, and slow the absorption of venom. Meat tenderizer (to neutralize the venom) or calamine lotion (to reduce itching) may be applied locally. If necessary, seek medical aid.
- In more serious reactions (severe and rapid swelling, allergic symptoms, and so forth) treat the bite/sting like you would treat a snakebite; that is, apply constricting bands above and below the site.
- Be prepared to perform basic life-supporting measures, such as rescue breathing.
- Reassure the casualty and keep him calm.
- In serious reactions, attempt to capture the insect for positive identification; however, be careful not to become a casualty yourself.
- If the reaction to the bite/sting appears serious, seek medical aid.

CAUTION

Insect bites/stings may cause *anaphylactic shock* (a shock caused by a severe allergic reaction). This is a life-threatening event and a **TRUE MEDICAL EMERGENCY**. Be prepared to perform the basic life-support measures and to immediately transport the casualty to an MTF.

NOTE

Be aware that some allergic or hypersensitive individuals may carry identification or emergency insect bite treatment kits. If the casualty is having an allergic reaction and has such a kit, administer the medication in the kit according to the instructions which accompany the kit.

d. Supplemental Information. For additional information concerning biting insects, see FM 8-230 and FM 21-10.

6-7. Table

See Table 6-1 for information on bites and stings.

Table 6-1. First Aid Measures for Bites and Stings

TYPES	FIRST AID MEASURES
Snakebite	<ol style="list-style-type: none"> 1. Move casualty away from the snake. 2. Remove jewelry from the affected area, if applicable. 3. Reassure casualty and keep him quiet. 4. Apply constricting band, 1-2 fingerbreadths proximal (near) to the bite. You should be able to insert a finger between the band and the skin. <ol style="list-style-type: none"> a. <i>Arm or Leg Bite.</i> Place one band above and one band below the bite site. b. <i>Hand or Foot Bite.</i> Place one band above the wrist or ankle. 5. Immobilize the affected part in a position below the level of the heart. 6. Kill the snake (if possible, without damaging its head or endangering yourself) and send it with the casualty. 7. Seek medical assistance immediately.
Brown Recluse Spider or Black Widow Spider Bite	<ol style="list-style-type: none"> 1. Keep casualty quiet. 2. Remove all jewelry from affected part, if applicable. 3. Wash the area. 4. Apply ice or freeze pack, if available. 5. Seek medical assistance.
Tarantula Bite or Scorpion Sting or Ant Bite	<ol style="list-style-type: none"> 1. Wash the area. 2. Remove all jewelry from the affected part, if applicable. 3. Apply ice or freeze pack, if available. 4. Apply baking soda, calamine lotion, or meat tenderizer (if available) to bite site to relieve pain and itching. 5. If the site of the bite is on the face, neck (possible airway problems), or genital area, or if local reaction seems severe, or if the sting is by the dangerous type of scorpion found in the Southwest United States desert, keep the casualty as quiet as possible. Seek medical assistance.

Bee Sting	<ol style="list-style-type: none">1. If the stinger is present, remove by scraping with a knife or fingernail. Do not squeeze venom sac on stinger; more venom may be injected.2. Remove all jewelry from affected part, if applicable.3. Wash the area.4. Apply ice or freeze pack, if available.5. If allergic signs or symptoms appear, be prepared to perform basic life support measures. Seek immediate medical assistance.
------------------	---

CHAPTER 7

FIRST AID IN A NUCLEAR, BIOLOGICAL, AND CHEMICAL ENVIRONMENT

7-1. General

American forces have not been exposed to NBC weapons/agents on the battlefield since World War I. In future conflicts and wars we can expect the use of such agents. Nuclear, biological, and chemical weapons will rapidly degrade unit effectiveness by forcing troops to wear protective clothing and by creating confusion and fear. Through training in protective procedures and first aid, units can maintain their effectiveness on the integrated battlefield.

7-2. First Aid Materials

You are issued materials to protect, decontaminate and use as first aid for NBC exposure. You must know how to use the items; some items are described in *a* through *c* below. It is equally important that you know when to use them.

a. Nerve Agent Pyridostigmine Pretreatment (NAPP). You will be issued a blister pack of pretreatment tablets when your commander directs. When ordered to take the pretreatment you must take one tablet every eight hours. This must be taken prior to exposure to nerve agents, since it may take several hours to develop adequate blood levels.

b. M291 Skin Decontaminating Kit. The M291 Skin Decontaminating Kit (Figure 7-1) contains six packets of XE-555 decontaminant resin.

WARNING

For external use only. May be slightly irritating to the eyes. Keep decontaminating powder out of eyes. Use water to wash toxic agent out of eyes. You may also use a 0.5 percent chlorine solution to wash toxic agent out of cuts or wounds.

c. Nerve Agent Antidote Kit, MARK I. Each soldier is issued three MARK I's for use in first aid for nerve agent poisoning (Figure 7-2).

Figure 7-1. M291 Skin Decontaminating Kit.

7-3. Classification of Chemical and Biological Agents

a. Chemical agents may be classified according to the primary physiological effects they produce, such as nerve, blister, blood, choking, vomiting, and incapacitating agents.

b. Biological warfare agents may be classified according to the effect they have on man. The effects include their ability to incapacitate, cause death, and how they are delivered. Most biological warfare agents are delivered as aerosols that effect the respiratory tract; some can be delivered by releasing infected insects, by contaminating food and water, and by injection (injecting material in individuals by terrorist, not mass exposure). These agents are found in living organisms such as fungi, bacteria, and viruses.

WARNING

Ingesting water or food contaminated with nerve, blister, and other chemical agents and with some biological agents can be fatal. NEVER consume water or food that is suspected of being contaminated until it has been tested and found safe for consumption by medical personnel.

7-4. Conditions for Masking Without Order or Alarm

a. Once an attack with a chemical or biological agent is detected or suspected, or information is available that such an agent is about to be used, you must STOP breathing and mask immediately. DO NOT WAIT to receive an order or alarm under the following circumstances:

- Your position is hit by artillery, that produce vapors, smoke, mist, etc or missiles, rockets, smokes, mists, and aerial in the offence of explosive devices.
- Smoke from an unknown source is present or approaching.
- A suspicious odor, liquid, or solid is present.
- A toxic chemical or biological attack is occurring.
- You are entering an area known or suspected of being contaminated.
- During any motor march, once chemical warfare has begun.
- When casualties are being received from an area where chemical or biological agents have reportedly been used.
- You have one or more of the following symptoms:

- An unexplained runny nose.
 - A sudden unexplained headache.
 - A feeling of choking or tightness in the chest or throat.
 - Dimness of vision.
 - Irritation of the eyes.
 - Difficulty in or increased rate of breathing without obvious reasons.
 - Sudden feeling of depression.
 - Dread, anxiety, restlessness.
 - Dizziness or light-headedness.
 - Slurred speech.
 - Unexplained laughter or unusual behavior is noted in others.
 - Numerous unexplained ill personnel.
 - Buddies suddenly collapsing without evident cause.
 - Animals or birds exhibiting unusual behavior or suddenly dying.
- b.* For further information, refer to FM 3-4, FM 8-10-7, FM 8-284, and FM 8-285.

7-5. First Aid for a Chemical Attack

Your field protective mask gives protection against biological and chemical agents as well as radiological fallout. With practice you can mask in 9 seconds or less, or put on your mask with hood within 15 seconds.

a. Stop breathing. Don your mask, seat it properly, and clear and check it; then resume breathing. Give the alarm, and continue the mission. Keep your mask on until the "all clear" signal has been given.

NOTE

Keep your mask on until the area is no longer hazardous and you are told to unmask.

b. If symptoms of nerve agent poisoning (paragraph 7-6) appear, immediately give yourself one MARK I.

CAUTION

Do not inject a nerve agent antidote until you are sure you need it.

c. If your eyes and face become contaminated, you must immediately try to get under cover. You need shelter to prevent further contamination while performing decontamination procedures your face. If no overhead cover is available put your poncho over your head before beginning the decontamination process. Then you put on the remaining protective clothing. If vomiting occurs, the mask should be lifted momentarily and drained--with your eyes are closed and while holding your breath and replaced, cleared, and sealed.

d. If nerve agents are used, mission permitting, watch for persons needing nerve agent antidotes and immediately follow procedures outlined in paragraph 7-8b.

e. When your mission permits, decontaminate your clothing and equipment.

7-6. Background Information on Nerve Agents

a. Nerve agents are among the deadliest of chemical agents. Nerve agents enter the body by inhalation, by ingestion, and through the skin. Depending on the route of entry and the amount, nerve agents can produce injury or death within minutes. Nerve agents can achieve their effects with small amounts. Nerve agents are absorbed rapidly, and the effects are felt immediately upon entry into the body. You will be issued three MARK I's and one Convulsant Antidote for Nerve Agent (CANA). Each MARK I consists of one atropine autoinjector and one pralidoxime chloride (2 PAM Cl) autoinjector. The CANA is a single autoinjector with flanges (Figure 7-2).

Figure 7-2. Nerve Agent Antidote Kit, MARK I and CANA.

b. When you have the signs and symptoms of nerve agent poisoning, you should immediately put on the protective mask and then inject yourself with one set of the MARK I. Do not administer the CANA. You should inject yourself in the outer (lateral) thigh muscle or if you are thin, in the upper outer (lateral) part of the buttocks.

c. Also, you may come upon an unconscious chemical agent casualty who will be unable to care for himself and who will require first aid. You should be able to successfully--

- (1) Mask him if he is unmasked.
- (2) Inject him, if necessary, with all of HIS autoinjectors.

- (3) Decontaminate his skin.
- (4) Seek medical aid.

7-7. Signs and Symptoms of Nerve Agent Poisoning

The symptoms of nerve agent poisoning are grouped as **MILD**--those that you recognize and for which you can perform self-aid, and **SEVERE**---those, which require buddy aid.

a. MILD Signs and Symptoms.

- Unexplained runny nose.
- Unexplained sudden headache.
- Sudden drooling.
- Difficulty seeing (dimness of vision and miosis).
- Tightness in the chest or difficulty in breathing.
- Localized sweating and muscular twitching in the area of contaminated skin.
- Stomach cramps.
- Nausea.
- Tachycardia followed by bradycardia. (*Tachycardia* is an abnormally rapid heart beat with a heart rate of over 100 beats per minute. *Bradycardia* is a slow heart rate of less than 60 beats per minute.)

b. SEVERE Signs and Symptoms.

- Strange or confused behavior.
- Wheezing, dyspnea (difficulty in breathing), and coughing.
- Severely pinpointed pupils.
- Red eyes with tearing.
- Vomiting.

- Severe muscular twitching and general weakness.
- Involuntary urination and defecation.
- Convulsions.
- Unconsciousness.
- Respiratory failure.
- Bradycardia.

7-8. First Aid for Nerve Agent Poisoning

First aid for nerve agent poisoning consists of administering the MARK I and CANA.

a. Injection Site. The injection site for administering the antidotes is normally in the outer thigh muscle. The thigh injection site is the area about a hand's width above the knee to a hand's width below the hip joint (Figure 7-3). It is important that the injection be given into a large muscle area. If the individual is thinly built, then the injections should be administered into the upper outer quarter (quadrant) of the buttock (Figure 7-4). Injecting in the buttocks of a thinly built individual avoids injury to the thighbone.

Figure 7-3. Thigh injection site.

Figure 7-4. Buttocks injection site.

b. Self-Aid. If you experience any or all of the nerve agent **MILD** symptoms (paragraph 7-7), you must **IMMEDIATELY** self-administer the MARK I (Figure 7-2). Follow the procedure given below.

- (1) Immediately put on your protective mask.
- (2) Remove one MARK I from your protective mask carrier, pocket of the MOPP overgarment, or other location as specified in your unit tactical standing operating procedure (TSOP). (In cold weather, the MARK I should be stored in an inside pocket of your clothing to protect the antidote from freezing. A frozen MARK I cannot be used to provide you with antidote, when needed. (However, the MARK I can still be used after complete thawing.)

(3) With your nondominant hand, hold the autoinjectors by the plastic clip so that the larger autoinjector is on top (Figure 7-5A) and both are positioned in front of you at eye level.

Figure 7-5. Removing the atropine autoinjector from the clip.

(4) With your dominant hand, check the injection site (thigh or buttock) for buttons or objects in pockets that may interfere with the injections.

(5) With the same hand, grasp the **atropine** autoinjector (the smaller of the two) with the thumb and first fingers (Figure 7-5B). **DO NOT** cover or hold the needle end with your hand, thumb, or fingers--you might accidentally inject yourself. An accidental injection into the hand **WILL NOT** deliver an effective dose of the antidote, especially if the needle goes through the hand.

(6) Pull the autoinjector out of the clip with a smooth motion (Figure 7-5C). The autoinjector is now armed. **DO NOT** touch the needle end.

(7) Hold the autoinjector with your thumb and two fingers (pencil writing position). Be careful not to inject yourself in the hand.

(8) Position the green (needle) end of the autoinjector against the injection site (thigh or buttock). **DO NOT** inject into areas close to the hip, knee, or thighbone.

(a) On the outer thigh muscle (Figure 7-6).

Figure 7-6. Thigh injection site for self-aid.

OR

(b) If you are thinly built, inject yourself into the upper outer quadrant of the buttock (Figure 7-7). There is a nerve that crosses the buttocks; hitting this nerve can cause paralysis. Therefore, you must only inject into the upper outer quadrant of the buttock.

Figure 7-7. Buttocks injection site for self-aid.

(9) Apply firm, even pressure (not a jabbing motion) to the autoinjector until it pushes the needle into your thigh (or buttock). Using a jabbing motion may result in an improper injection or injury to the thigh or buttock.

(10) Hold the autoinjector firmly in place for at least 10 seconds. The seconds can be estimated by counting "one thousand one," "one thousand two," and so forth. Firm pressure automatically triggers the coiled spring mechanism. This plunges the needle through the clothing into the muscle and at the same time injects the antidote into the muscle tissue.

(11) Carefully remove the autoinjector from the injection site.

(12) Place the used atropine autoinjector carefully between the little finger and the ring finger of the hand that is holding the remaining autoinjector and the clip (Figure 7-5A).

(13) Pull the **2 PAM Cl** autoinjector (the larger of the two) out of the clip (Figure 7-8B and C) and inject yourself in the same manner as steps (6) through (11) above, holding the black (needle) end against your outer thigh (or buttock). The same injection site can be used for both autoinjectors.

Figure 7-8. Removing the 2 PAM Cl autoinjector from the clip.

(14) Drop the clip without dropping the used autoinjectors.

(15) Attach the used autoinjectors to your clothing (Figure 7-9).

(a) Push the needles of the used autoinjectors (one at a time) through a pocket flap of your protective overgarment.

(b) Bend each needle to form a hook. It is important to keep track of all used autoinjectors so that medical personnel can determine how much antidote has already been given. Knowing how much antidote has been given enables them to provide the proper follow-up treatment, if needed. Be careful not to tear your protective garment and gloves with the needles.

Figure 7-9. One set of used autoinjectors attached to pocket flap.

(16) Massage the injection site, if time permits.

(17) After administering the first set of autoinjectors, you should decontaminate your skin, if necessary, and put on any remaining protective clothing.

(a) **DO NOT** give yourself another set of injections. If you are able to walk without assistance, know who you are, and where you are, you **WILL NOT** need the

second set of injections. (If not needed, giving yourself a second set of MARK I injections may create a nerve agent antidote overdose, which could cause incapacitation.)

(b) If you continue to have symptoms of nerve agent poisoning, seek someone else (a buddy) to check your symptoms and administer the additional sets of injections, if required.

c. *Buddy Aid.* Soldiers may seek assistance after self-aid (self-administering one MARK I) or may become incapacitated after self-aid. A buddy must evaluate the individual to determine if additional antidotes are required to counter the effects of the nerve agent. Also, soldiers may experience SEVERE symptoms of nerve agent poisoning (paragraph 7-7b); they will not be able to treat themselves. In either case, other soldiers must perform buddy aid as quickly as possible. Before initiating buddy aid, determine if one set of MARK I autoinjectors has already been used so that no more than three sets of the antidote are administered. Buddy aid also includes administering the CANA with the third MARK I to prevent convulsions. Follow the procedures indicated below.

WARNING

Squat, **DO NOT** kneel, when masking the casualty or administering the nerve agent antidotes to the casualty. Kneeling may force the chemical agent into or through your protective clothing.

- (1) Mask the casualty (follow the procedure outlined in FM 3-4).
- (2) Position the casualty on his side.
- (3) Position yourself near the casualty's thigh.
- (4) Remove all three MARK I sets (or the remaining sets) of the antidote autoinjectors and the CANA autoinjector from the inside pocket of the casualty's protective mask carrier, pocket of the MOPP suit, or other location specified in the unit TSOP.

CAUTION

DO NOT use your own MARK I's on a casualty. You may not have any antidote if needed for self-aid.

- (5) With your nondominant hand, hold the set of autoinjectors by the plastic clip, so that the larger autoinjector is on top (Figure 7-5A) and both are positioned in front of you at eye level.
- (6) With your dominant hand, check the injection site (thigh or buttock) for buttons or objects in pockets that may interfere with the injections.
- (7) With this same hand, grasp the atropine autoinjector (the smaller of the two) with the thumb and first fingers (Figure 7-5B). **DO NOT** cover or hold the needle end with your hand or fingers--you might accidentally inject yourself.

(8) Pull the autoinjector out of the clip with a smooth motion (Figure 7-5C). Make sure your hand **DOES NOT** cover the needle end. Holding or carrying the needle (green) end of the autoinjector may result in accidentally injecting yourself.

(9) Hold the autoinjector with your thumb and two fingers (pencil writing position). Be careful not to inject yourself in the hand.

(10) Position the green (needle) end of the autoinjector against the casualty's injection site (thigh or buttock).

(a) On the casualty's outer thigh muscle (Figure 7-10).

WARNING

DO NOT inject into areas close to the hip, knee, or thighbone.

Figure 7-10. Injecting the casualty's thigh.

OR

(b) If the casualty is thinly built, inject the antidote into the buttock. Only inject the antidote into the upper outer portion of the casualty's buttock (Figure 7-11). This avoids hitting the nerve that crosses the buttocks (Figure 7-4). Hitting this nerve can cause paralysis.

Figure 7-11. Injecting the casualty's buttocks.

(11) Apply firm, even pressure (not a jabbing motion) to the autoinjector until it pushes the needle into the casualty's thigh (or buttock). Make sure you do not hit the casualty's mask carrier or any objects in the individual's pockets. Using a jabbing motion may result in an improper injection or injury to the thigh or buttock.

(12) Hold the autoinjector firmly in place for at least 10 seconds. The seconds can be estimated by counting "one thousand one," "one thousand two," and so forth.

(13) Carefully remove the atropine autoinjector from the injection site.

(14) Place the used autoinjector carefully between the little finger and the ring finger of the hand that is holding the remaining autoinjector and the clip (Figure 7-8A).

(15) Pull the **2 PAM CI** autoinjector (the larger of the two) out of the clip (Figure 7-8B and C) and inject the casualty in the manner described in steps (7) through (12) above, holding the black (needle) end against the casualty's outer thigh (or buttock).

(16) Drop the clip *without dropping the used autoinjector*.

(17) Carefully lay the used autoinjectors on the casualty's side.

(18) Repeat the procedure immediately (steps [4] through [16] above), using the *second* and *third* sets of MARK I autoinjectors.

(19) Grasp the CANA autoinjector with your dominant hand with the needle end extending beyond your thumb and two fingers. With your other hand, pull the safety cap off the autoinjector base. The autoinjector is now armed. **DO NOT** touch the black (needle) end; to do so you may accidentally inject yourself.

(20) Position the black (needle) end of the autoinjector against the casualty's injection site (thigh or buttock).

(21) Apply firm, even pressure (not a jabbing motion) to the autoinjector until it pushes the needle into the casualty's thigh (or buttock). Make sure you do not hit the casualty's mask carrier or any objects in the individual's pockets.

(22) Hold the autoinjector firmly in place for at least 10 seconds. The seconds can be estimated by counting "one thousand one," "one thousand two," and so forth.

(23) Carefully remove the CANA autoinjector from the casualty's injection site.

(24) Attach the CANA and all three MARK I sets of used autoinjectors to the casualty's clothing (Figure 7-12).

(a) Push the needles of the used autoinjectors (one at a time) through one of the pocket flaps of the casualty's protective overgarment.

(b) Bend each needle to form a hook. Be careful **NOT** to tear the casualty's protective overgarments or your gloves with the needles.

(25) Massage the injection site if time permits.

Figure 7-12. Three sets of used MARK I autoinjectors and one CANA autoinjector attached to pocket flap.

c. *Combat Lifesaver.*

(1) The combat lifesaver must check to verify if the individual has received three sets of MARK I. If not, the combat lifesaver performs first aid as described for buddy aid above. If the individual has received the initial three sets of MARK I, then the combat lifesaver may administer additional atropine injections at approximately 15 minute intervals until atropinization is achieved (that is a heart rate above 90 beats per minute, reduced bronchial secretions, and reduced salivations). Administer additional atropine at intervals of 30 minutes to 4 hours to maintain atropinization or until the casualty is placed under the care of medical personnel. Check the heart rate by lifting the casualty's mask hood and feeling for a pulse at the carotid artery. Request medical assistance as soon as the tactical situation permits.

(2) The combat lifesaver should administer additional CANA to casualties suffering convulsions. Administer a second, and if needed, a third CANA at 5 to 10 minute intervals for a maximum of three injections (30 milligrams diazepam). Follow the steps and procedures described in buddy aid for administering the CANA. DO NOT give more than two additional injections for a total of three (one buddy aid plus two by the combat lifesaver).

7-9. Blister Agents

Blister agents (vesicants) include mustard (H and HD), nitrogen mustards (HN), lewisite (L), and other arsenicals, mixtures of mustards and arsenicals, and phosgene oxime (CX). Blister agents may act on the eyes, mucous membranes, lungs, and skin. They burn and blister the skin or any other body parts they contact. Even relatively low doses may cause serious injury. Blister agents damage the respiratory tract (nose, sinuses, and windpipe) when inhaled and cause vomiting and diarrhea when absorbed. Lewisite and phosgene oxime cause immediate pain on contact. However, mustard agents are deceptive as there is little or no pain at the time of exposure. Thus, in some cases, signs of injury may not appear for several hours after exposure.

a. Protective Measures. Your protective mask with hood and protective overgarment provide you protection against blister agents. If it is known or suspected that blister agents are being used, STOP BREATHING, put on your mask and all your protective overgarment.

CAUTION

Large drops of liquid vesicants on the protective overgarment ensemble may penetrate it if allowed to stand for an extended period. Remove large drops as soon as possible.

b. Signs and Symptoms of Blister Agent Poisoning.

(1) Immediate and intense pain upon contact with lewisite and phosgene oxime. No initial pain upon contact with mustard.

(2) Inflammation and blisters (burns) resulting in tissue destruction. The severity of a chemical burn is directly related to the concentration of the agent and the duration of

contact with the skin. The longer the agent is in contact with the tissue, the more serious the injury will be.

(3) Vomiting and diarrhea. Exposure to high concentrations of vesicants may cause vomiting or diarrhea.

(4) Death. The blister agent vapors absorbed during ordinary field exposure will probably not cause enough internal body (systemic) damage to result in death. However, death may occur from prolonged exposure to high concentrations of vapor or from extensive liquid contamination over wide areas of the skin, particularly *when decontamination is neglected or delayed*.

c. First Aid Measures.

(1) Use your M291 Skin Decontaminating Kit to decontaminate your skin and use water to flush contaminated eyes. Decontamination of vesicants must be done immediately (within 1 minute is best).

(2) If blisters form, cover them loosely with a field dressing and secure the dressing.

CAUTION

Blisters are actually burns. DO NOT attempt to decontaminate the skin where blisters have formed, as the agent has already been absorbed.

(3) If you receive blisters over a wide area of the body, you are considered seriously burned. Seek medical aid immediately.

(4) If vomiting occurs, the mask should be lifted momentarily and drained--while the eyes are closed and the breath is held--and replaced, cleared, and sealed.

(5) Remember, if vomiting or diarrhea occurs after having been exposed to blister agents, seek medical aid immediately.

7-10. Choking Agents (Lung-Damaging Agents)

Chemical agents that attack lung tissue, primarily causing fluid buildup (pulmonary edema), are classified as choking agents (lung-damaging agents). This group includes phosgene (CG), dipphosgene (DP), chlorine (Cl), and chloropicrin (PS). Of these four agents, phosgene is the most dangerous and is more likely to be employed by the enemy in future conflict.

a. Protective Measures. Your protective mask gives adequate protection against choking agents.

b. Signs and Symptoms. During and immediately after exposure to choking agents (depending on agent concentration and length of exposure), you may experience some or all of the following signs and symptoms:

- Tears (lacrimation).
- Coughing.
- Choking.
- Tightness of chest.
- Nausea and vomiting.
- Headaches.

c. Self-Aid.

(1) The protective mask should be put on immediately when any of the conditions described in *b* above exist. Another indication of a CG attack is an odor like newly mown hay. (Do not rely upon odor as indication of a chemical attack.)

(2) If some CG is inhaled, normal combat duties should be continued unless there is difficulty in breathing, nausea, vomiting, or more than the usual shortness of breath during exertion. If any of the above symptoms occur and the mission permits, remain at quiet rest until medical evacuation is accomplished.

d. Death. With ordinary field exposure to choking agents, death will probably not occur. *However, prolonged exposure to high concentrations of the vapor and neglect or delay in masking can be fatal.*

7-11. Blood Agents

Blood agents interfere with proper oxygen utilization in the body. Hydrogen cyanide (AC) and cyanogen chloride (CK) are the primary agents in this group.

a. Protective Measures. Your protective mask with a fresh filter gives adequate protection against field concentrations of blood agent vapor. The protective overgarment, as well as the mask, are needed when exposed to liquid AC.

b. Signs and Symptoms. During and immediately after exposure to blood agents (depending on agent concentration and length of exposure), you may experience some or all of the following signs and symptoms:

- Tearing (lacrimation).
- Eye, nose, and throat irritation.
- Sudden stimulation of breathing (unable to hold breath).
- Nausea.
- Coughing.
- Tightness of chest.
- Headache.
- Light-headedness (dizziness).
- Unconsciousness.

c. First Aid.

(1) *Hydrogen cyanide.* During any chemical attack, if you get a sudden stimulation of breath or detect an odor like bitter almonds, **PUT ON YOUR MASK IMMEDIATELY.** Speed is absolutely essential since this agent acts so rapidly that within a few seconds its effects will make it impossible for soldiers to put on their mask by themselves. Stop breathing until the mask is on, if at all possible. This may be very difficult since the agent strongly stimulates respiration.

(2) *Cyanogen chloride.* Put your mask on immediately if you experience any irritation of the eyes, nose, or throat. Soldiers who are unable to mask should be masked by the nearest soldier (buddy).

d. Medical Assistance. If you suspect that you have been exposed to blood agents, seek medical assistance immediately.

7-12. Incapacitating Agents

An incapacitating agent is a chemical agent which produces temporary, disabling conditions which persist for hours to days after exposure. Unlike riot control agents, which usually are momentary or fleeting in action, incapacitating agents have a persistent effect. It is likely that such agents will be disseminated by smoke-producing munitions or aerosols, thus making breathing their means of entry into the body. The protective mask is, therefore, essential.

a. There are no specific first aid measures to relieve the symptoms of incapacitating agents. Supportive first aid and physical restraint may be indicated. If the casualty is stuporous

or comatose, be sure that respiration is unobstructed; then turn him on his side in case vomiting should occur. Complete cleansing of the skin with soap and water should be done as soon as possible; or, the M291 Skin Decontaminating Kit can be used if washing is impossible. Remove weapons and other potentially harmful items from soldiers who are suspected of having these symptoms. Harmful items include cigarettes, matches, medications, and small items that might be swallowed accidentally. Delirious (confused) persons have been known to attempt to eat items bearing only a superficial resemblance to food.

b. Incapacitating agents (BZ type) may produce alarming dryness and coating of the lips and tongue; however, there is usually no danger of immediate dehydration. Fluids should be given sparingly, if at all, because of the danger of vomiting and because of the likelihood of temporary urinary retention due to paralysis of bladder muscles.

c. If the body temperature is elevated and mucous membranes are dry, immediate and vigorous cooling (as for heatstroke) is indicated. Methods that can be used to cool the skin are spraying with cool water and air circulation (fanning); applying alcohol soaked cloths and air circulation; and providing maximum exposure to air in a shaded area, along with maximum air circulation. Such cases are usually a result of anticholinergic poisoning. Rapid evacuation should be accomplished since medical treatment with the appropriate medication may be lifesaving.

CAUTION

Do not use ice for cooling the skin.

d. Reassurance and a firm, but friendly, attitude by individuals providing first aid will be beneficial if the casualty appears to comprehend what is being said. Conversation is a waste of time if the soldier is incoherent or cannot understand what is being said. In such cases, the less said, the better it is--these casualties will benefit more from prompt and vigorous restraint and evacuation to an MTF.

7-13. Incendiaries

Incendiaries can be grouped as white phosphorus, thickened gasoline, metal, and oil and metal. You must learn to protect yourself against these incendiaries.

a. White phosphorus is used primarily as a smoke producer but can be used for its incendiary effect to ignite field expedients and combustible materials. The burns from WP are usually multiple, deep, and variable in size. When particles of WP get on the skin or clothing, they continue to burn until deprived of air. They also have a tendency to stick to a surface and must be brushed off or picked out.

(1) If burning particles of WP strike and stick to your clothing, quickly take off the contaminated clothing before the WP burns through to the skin.

(2) If burning WP strikes your skin, smother the flame with water, a wet cloth, or mud.

NOTE

Since WP is soluble in oil, DO NOT use grease, oily ointments, or eye ointments to smother the flame.

(3) Keep the WP particles covered with a wet material to exclude air until you can remove them or have them removed from your skin.

(4) Remove the WP particles from the skin by brushing them with a wet cloth and by picking them out with a knife, bayonet, stick, or other available object.

(5) Seek medical aid once the mission permits.

b. Thickened fuel mixtures (napalm) have a tendency to cling to clothing and body surfaces, thereby producing prolonged exposure and severe burns. The first aid for these burns is the same as for other heat burns. The heat and irritating gases given off by these combustible mixtures may cause lung damage, which must be treated by medical personnel.

c. Metal incendiaries pose special problems. Thermite particles on the skin should be immediately cooled with water and then removed. The first aid for these burns is the same as for other heat burns. Particles of magnesium on the skin burn quickly and deeply. Like other metal incendiaries, they must be removed. Ordinarily, the complete removal of these particles should be done by medical personnel as soon as possible. Immediate medical treatment is required.

d. Oil and metal incendiaries have much the same effect on contact with the skin and clothing as those discussed (*b* and *c* above). First aid measures for burns are discussed in Chapter 3.

7-14. Biological Agents and First Aid

a. Biological attacks can result in combat ineffectiveness by introducing disease-causing organisms into a troop population.

b. Once a disease is identified, first aid or medical treatment is initiated depending on the seriousness of the disease. First aid measures are concerned with observable symptoms of the disease such as diarrhea or vomiting.

7-15. Toxins

Toxins are alleged to have been used in past conflicts. Witnesses and victims have described the agent as toxic rain (or yellow rain) because it was reported to have been released from aircraft as a yellow powder or liquid that covered the ground, structures, vegetation, and people.

a. Signs and Symptoms. The occurrence of the symptoms from toxins may appear in a period of a few minutes to several hours depending on the particular toxin, the soldier's susceptibility, and the amount of toxin inhaled, ingested, or deposited on the skin. Symptoms from toxins usually involve the central nervous system (CNS) but are often preceded by less prominent symptoms, such as nausea, vomiting, diarrhea, cramps, or stomach irritation and burning sensation. Typical neurological symptoms often develop rapidly in severe cases; for example, visual disturbances, inability to swallow, speech difficulty, lack of muscle coordination, and sensory abnormalities (numbness of mouth, throat, or extremities). Yellow rain (mycotoxins) also may have hemorrhagic symptoms which could include any or all of the following:

- Dizziness.
- Severe itching or tingling of the skin.
- Formation of multiple, small, hard blisters.
- Coughing up blood.
- Shock (which could result in death).

b. Self-Aid. Upon recognition of an attack employing toxins or the onset (start) of symptoms listed above, you must immediately take the following actions:

(1) Stop breathing, put on your protective mask with hood, and then resume breathing. Next, put on your protective clothing.

(2) Should severe itching of the face become unbearable, quickly—

- Loosen the cap on your canteen.
- Take and hold a deep breath and lift your mask.
- While holding your breath, close your eyes and flush your face with generous amounts of water.

CAUTION

DO NOT rub or scratch your eyes. Try not to let the water run onto your clothing or protective overgarment.

- Put your protective mask back on, seat it properly, clear it, and check it for seal; then resume breathing.
- Decontaminate your skin by bathing with soap and water as soon as the mission permits.
- Change clothing and decontaminate your protective mask using soap and water. Replace the filters if directed.

(3) If vomiting occurs, the mask should be lifted momentarily and drained--while the eyes are closed and the breath is held--and replaced, cleared, and sealed.

c. Medical Assistance. If you suspect that you have been exposed to toxins, you should seek medical assistance immediately.

7-16. Nuclear Detonation

a. Three types of injuries may result from a nuclear detonation. These are thermal, blast, and radiation injuries. Many times the casualty will have a combination of these types of injuries. First aid for thermal and blast injuries is provided based on observable injuries, such as burns, hemorrhage, or fractures. There is, however, no direct first aid for radiological casualties.

b. The signs and symptoms of radiation illness in the initial phase include the rapid onset of nausea, vomiting, and malaise (tiredness).

CHAPTER 8

FIRST AID FOR PSYCHOLOGICAL REACTIONS INTRODUCTION

8-1. General

Psychological first aid is as natural and reasonable as physical first aid and is just as familiar. When you were hurt as a child, the understanding attitude of your parents did as much as the psychological effect of a bandage. Later, your disappointment or grief was eased by supportive words from a friend. Certainly, taking a walk and talking things out with a friend are familiar ways of dealing with an emotional crisis. The same natural feelings that make us want to help a person who is injured make us want to give a helping hand to a buddy who is upset.

Psychological first aid really means nothing more complicated than assisting people with emotional distress whether it results from physical injury, disease, or excessive stress. Emotional distress is not always as visible as a wound or a broken bone. However, overexcitement, severe fear, excessive worry, deep depression, misdirected irritability, and anger are signs that stress has reached the point of interfering with effective coping. The more noticeable the symptoms become, the more urgent the need for you to be of help and the more important it is for you to know HOW to help.

8-2. Importance of Psychological First Aid

You must know how to give psychological first aid to be able to help yourself, your buddies, and your unit in order to keep performing the mission. Psychological first aid measures are simple and easy to understand. Your decision of what to do depends upon your ability to observe the soldier and understand his needs. Making the best use of resources requires ingenuity on your part. A stress reaction resulting in poor judgment can cause injury or even death to yourself or others on the battlefield. It can be even more dangerous if other persons are affected by the judgment of an emotionally upset soldier. If it is detected early enough, the affected soldier stands a good chance of remaining in his unit as an effective member. If it is not detected early and if the soldier becomes more emotionally upset, he may become a threat to himself and to others.

8-3. Situations Requiring Psychological First Aid

- Psychological first aid (buddy aid) is most needed at the first sign that a soldier cannot perform the mission because of emotional distress. Stress is inevitable in combat, in hostage and terrorist situations, and in civilian disasters such as floods, hurricanes, or industrial accidents. Most emotional reactions to such situations are temporary, and the soldier can still carry on with encouragement. Painful or disruptive symptoms may last for minutes, hours, or

days. However, if the stress symptoms are seriously disabling, they may be psychologically contagious and endanger not only the emotionally upset soldier but also the entire unit.

- Sometimes people continue to function well during a disastrous event, but suffer from emotional scars which impair their job performance or quality of life at a later time. Painful memories and dreams may recur for months and years and still be considered a normal reaction. However, if the memories are so painful that the person must avoid all situations which arouse them, becomes socially withdrawn, or shows symptoms of anxiety, depression, or substance abuse, they need treatment. Experience with police, firemen, emergency medical technicians, and others who deal with disasters has proved that the routine application of psychological first aid to all the participants, including those who have functioned well, greatly reduces the likelihood of future serious post-traumatic stress disorders (PTSDs).

8-4. Interrelationship of Psychological and Physical First Aid

Psychological first aid should go hand in hand with physical first aid. The discovery of a physical injury or cause for an inability to function does not rule out the possibility of a psychological injury (or vice versa). The person suffering from pain, shock, fear of serious injury, or fear of death does not respond well to joking, indifference, or fearful-tearful attention. Fear and anxiety may take as high a toll of the soldier's strength as does the loss of blood.

8-5. Goals of Psychological First Aid

The goals of psychological first aid are to—

- Be supportive; assist the soldier in dealing with his stress reaction.
- Prevent, and if necessary control, behavior harmful to himself and to others.
- Return the soldier to duty as soon as possible after dealing with the stress reaction.

8-6. Respect for Others' Feelings

a. Accept the soldier you are trying to help without censorship or ridicule. Respect his right to his own feelings. Even though your feelings, beliefs, and behavior are different, **DO NOT** blame or make light of him for the way he feels or acts. Your purpose is to help him in this tough situation, not to be his critic. A person **DOES NOT WANT** to be upset and worried. When he seeks help, he needs and expects consideration of his fears, not abrupt dismissal or ridicule.

b. Realize that people are the products of a wide variety of factors. All people **DO NOT** react the same way to the same situations. Each individual has complex needs and

motivations, both conscious and unconscious that are uniquely his own. Often the one thing that finally causes the person to become overloaded by a stressful situation is not the stressor itself, but some other problem.

8-7. Emotional and Physical Disability

a. Accept emotional disability as being just as real as physical disability. If a soldier's ankle is seriously sprained in a fall, no one expects him to run right away. A soldier's emotions may be temporarily strained by the overwhelming stress of battle or other traumatic incident. DO NOT demand that he pull himself together immediately and carry on without a break. Some individuals can pull themselves together immediately, but others cannot. The soldier whose emotional stability has been disrupted has a disability just as real as the soldier who has sprained his ankle. There is an unfortunate tendency in many people to regard as real only what they can see, such as a wound or bleeding. Some people tend to assume that damage involving a person's mind and emotions is just imagined, that he is not really sick or injured, and that he could overcome his trouble by using his will power.

b. The terms *it's all in your head*, *snap out of it*, and *get control of yourself* are often used by people who believe they are being helpful. Actually, these terms are expressions of hostility because they show lack of understanding. They only emphasize weakness and inadequacy. Such terms are of no use in psychological first aid.

c. Every physically injured person has some emotional reaction to the fact that he is injured.

(1) It is normal for an injured person to feel upset. The more severe the injury, the more insecure and fearful he becomes, especially if the injury is to a body part which is highly valued. For example, an injury to the eyes or the genitals, even though relatively minor, is likely to be extremely upsetting. An injury to some other part of the body may be especially disturbing to an individual for his own particular reason. For example, an injury of the hand may be a terrifying blow to a surgeon or an injury to the eye of a pilot.

(2) An injured soldier always feels less secure, more anxious, and more afraid not only because of what has happened to him but because of what he imagines may happen as a result of his injury. This fear and insecurity may cause him to be irritable, uncooperative, or unreasonable. As you help him, always keep in mind that such behavior has little or nothing to do with you personally. He needs your patience, reassurance, encouragement, and support.

8-8. Battle Fatigue and Other Combat Stress Reactions

Battle fatigue is a temporary emotional disorder or inability to function, experienced by a previously normal soldier as a reaction to the overwhelming or cumulative stress of combat. Battle fatigue gets better with reassurance, rest, physical replenishment, and activities that restore

confidence. All soldiers are likely to feel battle fatigue under conditions of intense and/or prolonged stress. They may even become battle fatigue casualties, unable to perform their mission for hours or days. Other combat stress reactions (CSRs) may result in negative behavior, but are not termed *battle fatigue* as they need more intensive treatment. These negative CSRs may result in misconduct stress behaviors such as drug and alcohol abuse, criminal acts, looting, desertion, and self-inflicted wounds. These harmful CSRs can often be prevented by good psychological first aid. Soldiers who commit misconduct stress behaviors may require disciplinary action rather than medical treatment.

8-9. Reactions to Stress

Most soldiers react to stressful incidents after the situation has passed. All soldiers feel some fear. This fear may be greater than they have experienced at any other time, or they may be more aware of their fear. In such a situation, they should not be surprised if they feel shaky or become sweaty, nauseated, or confused. These reactions are normal and are not a cause for concern. However, some reactions, either short- or long-term, will cause problems if left unchecked.

a. Emotional Reactions.

(1) The most CSR is inefficient performance. This can be demonstrated by--

- Slow thinking (or reaction time).
- Difficulty recognizing priorities and seeing what needs to be done.
- Difficulty getting started.
- Indecisiveness and having trouble focusing attention.
- Tendency to do familiar tasks and be preoccupied with familiar

details. (This can reach the point where the person is very passive, such as just sitting or wandering about not knowing what to do.)

(2) A less common reaction may be uncontrolled emotional outbursts; this can be demonstrated by crying, screaming, or laughing. Some soldiers will react in the opposite way. They will be very withdrawn and silent and try to isolate themselves from everyone. These soldiers should be encouraged to remain with their assigned unit. Uncontrolled reactions may appear by themselves or in any combination (the person may be crying uncontrollably one minute and then laughing the next). In this state, the person is restless and cannot keep still. He may run about, apparently without purpose. Inside, he feels a great rage or fear and his physical acts may show this. In his anger he may indiscriminately strike out at others.

b. Loss of Adaptability.

(1) In a desperate attempt to get away from the danger which has overwhelmed him, a soldier may panic and become confused. His mental ability may be so impaired he cannot think clearly or even follow simple commands. His judgement may be faulty and he may not be aware of his actions, such as standing up in his fighting position during an attack.

(2) In other cases, overwhelming stress may produce symptoms that are often associated with head injuries. For example, the soldier may appear dazed or be found wandering around aimlessly. He may appear confused and disoriented and may seem to have a complete or partial loss of memory. In such cases, especially when no eyewitnesses can provide evidence that the soldier has NOT suffered a head injury, it is necessary for him to be rapidly medically evacuated. **DO NOT ALLOW THE SOLDIER TO EXPOSE HIMSELF TO FURTHER PERSONAL DANGER UNTIL THE CAUSE OF THE PROBLEM HAS BEEN DETERMINED.**

c. Sleep Disturbance and Repetition of Dreams. A person who has been overwhelmed by stress often has difficulty sleeping. The soldier may experience nightmares related to the stressors. Remember that nightmares, in themselves, are not considered abnormal when they occur soon after a period of intensive stress. As time passes, the nightmares usually become less frequent and less intense. In extreme cases, a soldier, even when awake, may think repeatedly of the incident, feel as though it is happening again, and act out parts of his stress over and over again. For some persons, this repetitious reexperiencing of the stressful event may be necessary for eventual recovery; therefore, it should not be discouraged or viewed as abnormal. For the person reexperiencing the event, such reaction may be disruptive. The soldier needs to be encouraged to *ventilate* about the incident. Ventilation is a technique where the soldier is given the opportunity to talk extensively, often repetitiously about the experience.

8-10. Severe Stress or Battle Fatigue

You do not need specialized training to recognize severe stress or battle fatigue that will cause problems for the soldier, the unit, or the mission. Reactions that are less severe, however, are more difficult to detect. To determine whether a person needs help, you must observe him to see whether he is doing something meaningful, performing his duties, taking care of himself, behaving in an unusual fashion, or acting out of character.

8-11. Application of Psychological First Aid

The emotionally disturbed soldier has built a barrier against fear. He does this for his own protection, although he is probably not aware that he is doing it. If he finds that he does not have to be afraid and that there are normal, understandable things about him, he will feel safer in dropping this barrier. Persistent efforts to make him realize that you want to understand him will be reassuring, especially if you remain calm. Nothing can cause an emotionally disturbed person to become even more fearful than feeling that others are afraid of him. Try to remain calm.

Familiar things, such as a cup of coffee, the use of his name, attention to a minor wound, being given a simple job to do, or the sight of familiar people and activities, will add to his ability to overcome his fear. He may not respond well if you get excited, angry, or abrupt.

a. Ventilation. After the soldier becomes calmer, he is likely to have dreams about the stressful event. He also may think about it when he is awake or even repeat his personal reaction to the event. One benefit of this natural pattern is that it helps him master the stress by going over it just as one masters the initial fear of parachuting from an aircraft by doing it over and over again. Eventually, it is difficult to remember how frightening the event was initially. In giving first aid to the emotionally disturbed soldier, you should let him follow this natural pattern. Encourage him to talk. Be a good listener. Let him tell, in his own words, what actually happened. If home front problems or worries have contributed to the stress, it will help him to talk about them. Your patient listening will prove to him that you are interested in him, and by describing his personal problem, he can work at mastering his fear. If he becomes overwhelmed in the telling, suggest a cup of coffee or a break. Whatever you do, assure him that you will listen again as soon as he is ready. Do try to help put the soldier's perception of what happened back into realistic perspective; but, DO NOT argue about it.

b. Activity.

(1) A person who is emotionally disturbed as the result of a combat action is a casualty of anxiety and fear. He is disabled because he has become temporarily overwhelmed by his anxiety. A good way to control fear is through activity. Almost all soldiers, for example, experience a considerable sense of anxiety and fear while they are poised, awaiting the opening of a big offensive; but this is normally relieved, and they actually feel better once they begin to move into action. They take pride in effective performance and pleasure in knowing that they are good soldiers, perhaps being completely unaware that overcoming their initial fear was their first major accomplishment.

(2) Useful activity is very beneficial to the emotionally disturbed soldier who is not physically incapacitated. After you help a soldier get over his initial fear, help him to regain some self-confidence. Make him realize his job is continuing by finding him something useful to do. Encourage him to be active. Get him to help load trucks, clean up debris, or dig fighting positions. If possible, get him back to his usual duty. Seek out his strong points and help him apply them. Avoid having him just sit around. You may have to provide direction by telling him what to do and where to do it. The instructions should be clear and simple and should be repeated. A person who has panicked is likely to argue. Respect his feelings, but point out more immediate, obtainable, and demanding needs. Channel his excessive energy and, above all, DO NOT argue. If you cannot get him interested in doing more profitable work, it may be necessary to enlist aid in controlling his overactivity before it spreads to the group and results in more panic. Prevent the spread of such infectious feelings by restraining and segregating if necessary.

(3) Involvement in activity helps a soldier in three ways; he--

- Forgets himself.

- Has an outlet for his excessive tensions.
- Proves to himself he can do something useful.

c. Rest. There are times, particularly in combat, when physical exhaustion is a principal cause for emotional reactions. A unit sleep plan should be established and implemented. When possible, soldiers should be given a safe and relatively comfortable area in which to sleep. Examples would be an area away from heavy traffic, noise, and congestion or a place that is clean and dry and protected from environmental conditions. The more uninterrupted sleep a soldier gets the better he will be able to function in the tactical environment.

d. Hygiene. Field hygiene is an important ingredient in a soldier's morale. A soldier who is dirty and unkempt will not function as well as a soldier who has had the opportunity to bathe and put on clean, dry clothing. During combat, unit leaders should stress the importance of personal hygiene. Good personal hygiene not only improves morale, it also is a preventive measure against disease and nonbattle injury (DNBI).

e. Group Activity. You have probably already noticed that a person works, faces danger, and handles serious problems better if he is a member of a closely-knit group. Each soldier in the team supports the other team members. Esprit de corps is built because the soldiers have the same interests, goals, and mission, and as a result they are more productive; furthermore, they are less worried because everyone is involved. It is this spirit that takes a strategic hill in battle. It is so powerful that it is one of the most effective tools you have in your *psychological first aid bag*. Getting the soldier back into the team or squad activities will reestablish his sense of belonging and security and will go far toward making him a useful member of the unit.

8-12. Reactions and Limitations

Up to this point the discussion has been primarily about the feelings of the emotionally distressed soldier. What about your feelings toward him? Whatever the situation, you will have emotional reactions (conscious or unconscious) toward this soldier. Your reactions can either help or hinder your ability to help him. When you are tired or worried, you may very easily become impatient with him if he is unusually slow or exaggerates. You may even feel resentful toward him. At times when many physically wounded lie about you, it will be especially natural for you to resent disabilities that you cannot see. Physical wounds can be seen and easily accepted. Emotional reactions are more difficult to accept as injuries. On the other hand, will you tend to be overly sympathetic? Excessive sympathy for an incapacitated person can be as harmful as negative feelings in your relationship with him. He needs strong help, but not your sorrow. To overwhelm him with pity will make him feel even more inadequate. You must expect your buddy to recover, to be able to return to duty, and to become a useful soldier again. This expectation should be displayed in your behavior and attitude as well as in what you say. If he

can see your calmness, confidence, and competence, he will be reassured and will feel a sense of greater security.

8-13. Tables.

See Tables 8-1, 8-2, and 8-3 for more information.

Table 8-1. Mild Battle Fatigue

<u>PHYSICAL SIGNS*</u>	<u>EMOTIONAL SIGNS*</u>
1. Trembling, tearful	1. Anxiety, indecisiveness
2. Jumpiness, nervousness	2. Irritable, complaining
3. Cold sweat, dry mouth	3. Forgetful, unable to concentrate
4. Pounding heart, dizziness	4. Insomnia, nightmares
5. Nausea, vomiting, diarrhea	5. Easily startled by noises, movement
6. Fatigue	6. Grief, tearful
7. <i>Thousand-yard stare.</i>	7. Anger, beginning to lose confidence in self and unit
8. Difficulty thinking, speaking, and communicating	

SELF- AND BUDDY AID

1. Continue mission performance, focus on immediate mission.
2. Expect soldier to perform assigned duties.
3. Remain calm at all times; be directive and in control.
4. Let soldier know his reaction is normal, and that there is nothing seriously wrong with him.
5. Keep soldier informed of the situation, objectives, expectations, and support. Control rumors.
6. Build soldier's confidence, talk about succeeding.
7. Keep soldier productive (when not resting) through recreational activities, equipment maintenance.
8. Ensure soldier maintains good personal hygiene.
9. Ensure soldier eats, drinks, and sleeps as soon as possible.
10. Let soldier talk about his feelings. DO NOT "put down" his feelings of grief or worry. Give practical advice and put emotions into perspective.

*Most or all of these signs are present in mild battle fatigue. They can be present in any normal soldier in combat yet he can still do his job.

Table 8-2. *More Serious Battle Fatigue*

<u>PHYSICAL SIGNS*</u>		<u>EMOTIONAL SIGNS*</u>	
1.	Constantly moves around	1.	Rapid and/or inappropriate talking
2.	Flinching or ducking at sudden sounds	2.	Argumentative, reckless movements/actions.
3.	Shaking, trembling (whole body or arms)	3.	Inattentive to personal hygiene.
4.	Cannot use part of body, no physical reason (hand, arm, legs)	4.	Indifferent to danger
5.	Cannot see, hear, or feel (partial or complete loss)	5.	Memory loss
6.	Physical exhaustion, crying	6.	Severe stuttering, mumbling, or cannot speak at all
7.	Freezing under fire, or total immobility	7.	Insomnia, nightmares
8.	Vacant stares, staggers, sways when stands	8.	Seeing or hearing things that do not exist
9.	Panic running under fire	9.	Rapid emotional shifts
		10.	Social withdrawal
		11.	Apathetic
		12.	Hysterical outbursts
		13.	Frantic or strange behavior

TREATMENT PROCEDURES**

1. If a soldier's behavior endangers the mission, self, or others, do whatever necessary to control the soldier.
2. If the soldier is upset, calmly talk him into cooperating.
3. If concerned about the soldier's reliability:
 - Unload the soldier's weapon.
 - Take weapon if seriously concerned.
 - Physically restrain the soldier only when necessary for safety or transportation.
4. Reassure everyone that the signs are probably just battle fatigue and will quickly improve.
5. If battle fatigue signs continue:
 - Get the soldier to a safer place.
 - DO NOT leave the soldier alone, keep someone he knows with him.
 - Notify senior NCO or officer.
 - Have the soldier examined by medical personnel.
6. Give the soldier easy tasks to do when not sleeping, eating, or resting.
7. Assure the soldier he will return to full duty in 24 hours; and, return soldier to normal duties as soon as he is ready.

*These signs are present in addition to the signs of mild BF reaction.

**Do these procedures in addition to the self and buddy aid care.

Table 8-3. Preventive Measures to Combat Battle Fatigue

1. Welcome new members into your team, get to know them quickly. If you are new, be active in making friends.
2. Be physically fit (strength, endurance, and agility).
3. Know and practice lifesaving self and buddy aid.
4. Practice rapid relaxation techniques (FM 22-51).
5. Help each other out when things are tough at home or in the unit.
6. Keep informed; ask your leader questions, ignore rumors.
7. Work together to give everyone food, water, shelter, hygiene, and sanitation.
8. Sleep when mission and safety permit; let everyone get time to sleep.
 - Sleep only in safe places and by SOP.
 - If possible, sleep 6 to 9 hours per day.
 - Try to get at least 4 hours sleep per day.
 - Get good sleep before going on sustained operations.
 - Catnap when you can, but allow time to wake up fully.
 - Catch up on sleep after going without.

APPENDIX A

FIRST AID CASE, KITS, DRESSINGS, AND BANDAGES

A-1. First Aid Case with Field Dressings and Bandages

Every *service member* is issued a first aid case (Figure A-1A) with a field first aid dressing encased in a plastic wrapper (Figure A-1B). He carries it at all times for his use. The field first aid dressing is a standard sterile (germ-free) compress or pad with bandages attached (Figure A-1C). This dressing is used to cover the wound, to protect against further contamination, and to stop bleeding (pressure dressing). *When a service member administers first aid to another person, he must remember to use the wounded person's dressing; he may need his own later.* The *service member* must check his first aid case regularly and replace any used or missing dressing. The field first aid dressing may normally be obtained from his unit supply.




Figure A-1. Field first aid case and dressing (Illustrated A through C).

A-2. General Purpose First Aid Kits

General purpose first aid kits for Army personnel are listed in the common table of allowances (CTA) 8-100. These kits are carried on Army vehicles, aircraft, and boats for use by the operators, crew, and passengers. Individuals designated by unit TSOP to be responsible for these kits are required to check them regularly and replace all items used. The general purpose kit and its contents can be obtained through the unit supply system.

NOTE

Periodically check the dressings (for holes or tears in the packaging) and the medicines (for expiration date) that are in the first aid kits. If necessary, replace defective or outdated items.

A-3. Dressings

1
2 Dressings are *sterile pads* or *compresses* used to cover wounds. They usually are made of gauze
3 or cotton wrapped in gauze (Figure A-1C). In addition to the standard field first aid dressing,
4 other dressings, such as sterile gauze compresses and small sterile compresses on adhesive strips,
5 may be available under CTA 8-100.

6
7

8 **A-4. Standard Bandages**

9

10 *a.* Standard bandages are made of gauze or muslin and are used over a sterile
11 dressing to secure the dressing in place, to close off its edge from dirt and germs, and to create
12 pressure on the wound and control bleeding. A bandage can also support an injured part or
13 secure a splint.

14

15 *b.* Tailed bandages may be attached to the dressing as indicated on the field first aid
16 dressing (Figure A-1C).

17

18

19 **A-5. Triangular and Cravat (Swathe) Bandages**

20

21 *a.* Triangular and cravat (or swathe) bandages (Figure A-2) are fashioned from a
22 triangular piece of muslin (37 x 37 x 52 inches) provided in the general purpose first aid kit. If it
23 is folded into a strip, it is called a cravat. Two safety pins are packaged with each bandage.
24 These bandages are valuable in an emergency since they are easily applied.


25

26 *b.* To improvise a triangular bandage, cut a square of available material, slightly
27 larger than 3 feet by 3 feet, and **FOLD** it **DIAGONALLY**. If two bandages are needed, cut the
28 material along the **DIAGONAL FOLD**.

29

30 *c.* A cravat can be improvised from such common items as T-shirts, other shirts, bed
31 linens, trouser legs, scarfs, or any other item made of pliable and durable material that can be
32 folded, torn, or cut to the desired size.

33



1
2

Figure A-2. Triangular and cravat bandages (Illustrated A through E).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

APPENDIX B

RESCUE AND TRANSPORTATION PROCEDURES

B-1. General

A basic principle of first aid is to evaluate the casualty's injuries and administer first aid before moving him. However, adverse situations or conditions may jeopardize the lives of both the rescuer and the casualty if this is done. It may be necessary *first to rescue* the casualty before first aid can be effectively or safely given. The life and/or the well-being of the casualty will depend as much upon the manner in which he is *rescued and transported*, as it will upon the first aid and medical treatment he receives. Rescue actions must be done quickly and safely. Careless or rough handling of the casualty during rescue operations can aggravate his injuries.

B-2. Principles of Rescue Operations

a. When faced with the necessity of rescuing a casualty who is threatened by hostile action, fire, water, or any other immediate hazard, **DO NOT** take action without first determining the extent of the hazard and your ability to handle the situation. **DO NOT** become a casualty.

b. The rescuer must evaluate the situation and analyze the factors involved. This evaluation involves three major steps:

- Identify the task.
- Evaluate circumstances of the rescue.
- Plan the action.

B-3. Considerations

a. First determine if a rescue attempt is actually needed. It is a waste of time, equipment, and personnel to rescue someone not in need of rescuing. It is also a waste to look for someone who is not lost or needlessly risk the lives of the rescuer(s). In planning a rescue, attempt to obtain the following information:

- Who, what, where, when, why, and how the situation happened?
- How many casualties are involved and the nature of their injuries?
- What is the tactical situation?

- 1 • What are the terrain features and the location of the casualties?
- 2
- 3 • Will there be adequate assistance available to aid in the rescue/evacuation?
- 4
- 5 • Can first aid and/or medical treatment be provided at the scene; will the
- 6 casualties require movement to a safer location?
- 7
- 8 • What specialized equipment will be required for the rescue operation?
- 9
- 10 • Is the rescue area contaminated? Will decontamination equipment and
- 11 materiel be required for casualties, rescue personnel, and rescue equipment?
- 12
- 13 • How much time is available?
- 14

15 *b.* The time element can play a significant role in how the rescue is attempted. If the
16 casualties are in imminent danger of losing their lives (such as near a burning vehicle or in a
17 burning building), the time available will be relatively short and will sometimes cause a rescuer
18 to compromise planning stages and/or the first aid which can be given. However, if the casualty
19 is in a relatively secure area and his physical condition is strong, more deliberate planning can
20 take place. A realistic estimate of time available must be made as quickly as possible to
21 determine action time remaining. The key elements are the casualty's physical and mental
22 condition, the tactical situation, and the environment.

23

24

25 **B-4. Plan of Action**

26

27 *a.* The casualty's ability to endure is of primary importance in estimating the time
28 available. Age, physical condition, and extent of wounds and/or injuries will differ from
29 casualty to casualty. Therefore, to determine the time available, you will have to consider—

- 30
- 31 • Extent of injuries.
- 32
- 33 • Type of situation.
- 34
- 35 • Personnel and/or equipment availability.
- 36
- 37 • Weather.
- 38
- 39 • Terrain (natural and man-made).
- 40
- 41 • Environment (contaminated or uncontaminated).
- 42

43 *b.* In respect to terrain, you must consider altitude and visibility. In some cases, the
44 casualty may be of assistance because he knows more about the particular terrain or situation
45 than you do. Maximum use of secure/reliable trails or roads is essential.

1 c. When taking weather into account, ensure that blankets and/or rain gear are
2 available. Even a mild rain can complicate a normally simple rescue. In high altitudes and/or
3 extreme cold and gusting winds, the time available is critically shortened. Be prepared to
4 provide shelter and warmth for the casualty as well as the rescuers.
5
6

7 **B-5. Proper Handling of Casualties**

8

9 a. You may have saved the casualty's life through the application of appropriate first
10 aid measures. However, his life can be lost through rough handling or careless transportation
11 procedures. Before you attempt to move the casualty—
12

- 13 • Evaluate the type and extent of his injuries.
- 14 • Ensure that dressings over wounds are adequately reinforced.
- 15 • Ensure that fractured bones are properly immobilized and supported to
16 prevent them from cutting through muscle, blood vessels, and skin.
17
18

19 b. Based upon your evaluation of the type and extent of the casualty's injury and
20 your knowledge of the various manual carries, you must select the best possible method of
21 manual transportation. If the casualty is conscious, tell him how he is to be transported. This
22 will help allay his fear of movement and gain his cooperation and confidence.
23
24

25 c. Buddy aid for chemical agent casualties includes those actions required to prevent
26 an incapacitated casualty from receiving additional injury from the effects of chemical hazards.
27 If a casualty is physically unable to decontaminate himself or administer the proper chemical
28 agent antidote, the casualty's buddy assists him and assumes responsibility for his care. Buddy-
29 aid includes—
30

- 31 • Administering the proper chemical agent antidote.
- 32 • Decontaminating the incapacitated casualty's exposed skin.
- 33 • Ensuring that his protective ensemble remains correctly emplaced.
- 34 • Maintaining respiration.
- 35 • Controlling bleeding.
- 36 • Providing other standard first aid measures.
- 37 • Transporting the casualty out of the contaminated area.
38
39
40
41
42
43
44
45
46

1 **B-6. Positioning the Casualty**
2

3 The first step in any manual carry is to position the casualty to be lifted. If he is conscious, he
4 should be told how he is to be positioned and transported. This helps lessen his fear of
5 movement and to gain his cooperation. It may be necessary to roll the casualty onto his
6 abdomen, or his back, depending upon the position in which he is lying and the particular carry
7 to be used.
8

9 a. To roll a casualty onto his abdomen, kneel at the casualty's uninjured side.

10
11 (1) Place his arms above his head; cross his ankle which is farther from you
12 over the one that is closer to you.

13
14 (2) Place your hands on the shoulder which is farther from you; place your
15 other hand in the area of his hip or thigh (Figure B-1).

16
17 (3) Roll him gently toward you onto his abdomen (Figure B-2).
18



19
20 *Figure B-1. Positioning the casualty.*
21



22
23 *Figure B-2. Rolling casualty onto his abdomen.*
24
25

1 *b.* To roll a casualty onto his back, follow the same procedure described in *a* above,
2 except gently roll the casualty onto his back, rather than onto his abdomen.
3
4

5 **B-7. Medical Evacuation and the Transportation of Casualties**

6

7 *a.* Medical evacuation of the sick and wounded (with en route medical care) is the
8 responsibility of medical personnel who have been provided special training and equipment.
9 Therefore, unless a good reason for you to transport a casualty arises, wait for some means of
10 medical evacuation to be provided. When the situation is urgent and you are unable to obtain
11 medical assistance or know that no medical evacuation assets are available, you will have to
12 transport the casualty. For this reason, you must know how to transport him without increasing
13 the seriousness of his condition.
14

15 *b.* Transporting a casualty by litter (FM 8-10-6) is safer and more comfortable for
16 him than by manual means; it is also easier for you. Manual transportation, however, may be the
17 only feasible method because of the terrain or the combat situation; or it may be necessary to
18 save a life. In these situations, the casualty should be transferred to a litter as soon as one can be
19 made available or improvised.
20

21 **B-8. Manual Carries**

22

23 Casualties carried by manual means must be carefully and correctly handled, otherwise their
24 injuries may become more serious or possibly fatal. Situation permitting, transport of a casualty
25 should be organized and unhurried. Each movement should be performed as deliberately and
26 gently as possible. Casualties should not be moved before the type and extent of injuries are
27 evaluated and the required first aid is administered. The exception to this occurs when the
28 situation dictates immediate movement for safety purposes (for example, it may be necessary to
29 remove a casualty from a burning vehicle); that is, the situation dictates that the urgency of
30 casualty movement outweighs the need to administer first aid. Manual carries are tiring for the
31 bearers and involve the risk of increasing the severity of the casualty's injury. In some instances,
32 however, they are essential to save the casualty's life. Although manual carries are accomplished
33 by one or two bearers, the two-man carries are used whenever possible. They provide more
34 comfort to the casualty, are less likely to aggravate his injuries, and are also less tiring for the
35 bearers. The distance a casualty can be carried depends on many factors, such as—
36

- 37 • Nature of the casualty's injuries.
- 38
- 39 • Strength and endurance of the bearer(s).
- 40
- 41 • Weight of the casualty.
- 42
- 43 • Obstacles encountered during transport (natural or man-made).
- 44
- 45 • Type of terrain.
- 46

1 *a. One-Man Carries.* These carries should be used when only one bearer is
2 available to transport the casualty.

3
4 (1) The *fireman's carry* (Figure B-3) is one of the easiest ways for one
5 individual to carry another. After an unconscious or disabled casualty has been properly
6 positioned, he is raised from the ground, then supported and placed in the carrying position.

7
8 (*a*) After rolling the casualty onto his abdomen, straddle him. Extend
9 your hands under his chest and lock them together.

10
11 (*b*) Lift the casualty to his knees as you move backward.

12
13 (*c*) Continue to move backward, thus straightening the casualty's legs
14 and locking his knees.

15
16 (*d*) Walk forward, bringing the casualty to a standing position; tilt him
17 slightly backward to prevent his knees from buckling.

18
19 (*e*) As you maintain constant support of the casualty with one arm,
20 free your other arm, quickly grasp his wrist, and raise his arm high. Instantly pass your head
21 under his raised arm, releasing it as you pass under it.


22
23 (*f*) Move swiftly to face the casualty and secure your arms around his
24 waist. Immediately place your foot between his feet and spread them apart (approximately 6 to 8
25 inches).

26
27 (*g*) Grasp the casualty's wrist and raise his arm high over your head.

28
29 (*h*) Bend down and pull the casualty's arm over and down on your
30 shoulder, bringing his body across your shoulders. At the same time, pass your arm between his
31 legs.


32
33 (*i*) Grasp the casualty's wrist with one hand, and place your other
34 hand on your knee for support.

35
36 (*j*) Rise with the casualty positioned correctly. Your other hand is
37 free for use.



1
2

Figure B-3. Fireman's carry.



3
4

Figure B-3. Fireman's carry (continued).

1




Figure B-3. Fireman's carry (continued).

2
3
4
5
6
7
8
9
10
11
12
13
14

(2) The alternate method of the *fireman's carry* for raising a casualty from the ground is illustrated in Figure B-4; however, it should be used only when the bearer believes it to be safer for the casualty because of the location of his wounds. When the alternate method is used, care must be taken to prevent the casualty's head from snapping back and causing a neck injury. The steps for raising a casualty from the ground for the fireman's carry are also used in other one-man carries.

(a) Kneel on one knee at the casualty's head and face his feet. Extend your hands under his armpits, down his sides, and across his back.

1
2
3
4

(b) As you rise, lift the casualty to his knees. Then secure a lower hold and raise him to a standing position with his knees locked.



5
6
7
8
9

Figure B-4. Fireman's carry (alternate method) for lifting a casualty to a standing position.

10
11
12
13
14
15
16
17
18
19
20
21

(3) In the *supporting carry* (Figure B-5), the casualty must be able to walk or at least hop on one leg, using the bearer as a crutch. This carry can be used to assist him as far as he is able to walk or hop.

(a) Raise the casualty from the ground to a standing position by using the fireman's carry.

(b) Grasp the casualty's wrist and draw his arm around your neck.

(c) Place your arm around his waist. The casualty is now able to walk or hop using you as a support.



Figure B-5. Supporting carry.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17

(4) The *arms carry* (Figure B-6) is useful in carrying a casualty for a short distance (up to 50 meters) and for placing him on a litter.

(a) Raise or lift the casualty from the ground to a standing position, as in the fireman's carry.

(b) Place one arm under the casualty's knees and your other arm around his back.

(c) Lift the casualty.

(d) Carry the casualty high to lessen fatigue.



Figure B-6. Arms carry.

1
2
3
4
5 (5) Only a conscious casualty can be transported by the *saddleback carry*
6 (Figure B-7), because he must be able to hold onto the bearer's neck. To use this technique—
7

8 (a) Raise the casualty to an upright position, as in the fireman's carry.
9

10 (b) Support the casualty by placing an arm around his waist. Move to
11 the casualty's side. Have the casualty put his arm around your neck and move in front of him
12 with your back to support him.
13

14 (c) Have the casualty encircle his arms around your neck
15

16 (d) Stoop, raise him on your back and clasp your hands together
17 beneath his thighs, if possible.
18



Figure B-7. Saddleback carry.

1
2
3
4
5 (6) In the *pack-strap carry* (Figure B-8), the casualty's weight rests high on
6 the your back. This makes it easier for you to carry the casualty a moderate distance (50 to 300
7 meters). To eliminate the possibility of injury to the casualty's arms, you must hold his arms in a
8 palms-down position.
9

10 (a) Lift the casualty from the ground to a standing position, as in the
11 fireman's carry.

12
13 (b) Support the casualty with your arms around him and grasp his
14 wrist closer to you.

15
16 (c) Place his arm over your head and across your shoulders.

17
18 (d) Move in front of him while still supporting his weight against your
19 back.

20
21 (e) Grasp his other wrist and place this arm over your shoulder.

22
23 (f) Bend forward and raise or hoist the casualty as high on your back
24 as possible so that his weight is resting on your back.

1
2
3
4
5

NOTE

Once the casualty is positioned on the bearer's back, the bearer remains as erect as possible to prevent straining or injuring his back.



Figure B-8. Pack-strap carry.

6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30

(7) The *pistol-belt carry* (Figure B-9) is the best one-man carry for a long distance (over 300 meters). The casualty is securely supported upon your shoulders by a belt. Both your hands and the casualty's (if conscious) are free for carrying a weapon or equipment, climbing obstacles. With your hands free and the casualty secured in place, you are also able to creep through shrubs and under low-hanging branches.

(a) Link two pistol belts (or three, if necessary) together to form a sling. Place the sling under the casualty's thighs and lower back so that a loop extends from each side.

NOTE

If pistol belts are not available for use, other items such as a rifle sling, two cravat bandages, two litter straps, or any other suitable material that will not cut or bind the casualty may be used.

(b) Lie face up between the casualty's outstretched legs. Thrust your arms through the loops and grasp his hands and trouser leg on his injured side.

(c) Roll toward the casualty's uninjured side onto your abdomen, bringing him onto your back. Adjust the sling, if necessary.

1 (d) Rise to a kneeling position. The belt will hold the casualty in
2 place.

3
4 (e) Place one hand on your knee for support and rise to an upright
5 position. (The casualty is supported on your shoulders.)

6
7 (f) Carry the casualty with your hands free for use in rifle firing,
8 climbing, or surmounting obstacles.

9




Figure B-9. Pistol-belt carry.

10
11
12
13
14
15
16
17

(8) The *pistol-belt drag* (Figure B-10), as well as other drags, is generally used for short distances (up to 50 meters). This drag is useful in combat, since both the bearer and the casualty can remain closer to the ground than in any other drags.

1 (a) Extend two pistol belts or similar objects to their full length and
2 join them together to make a continuous loop.

3
4 (b) Roll the casualty onto his back, as in the fireman's carry.

5
6 (c) Pass the loop over the casualty's head, and position it across his
7 chest and under his armpits. Then cross the remaining portion of the loop, thus forming a figure
8 eight.

9
10 (d) Lie on your side facing the casualty.

11
12 (e) Slip the loop over your head and turn onto your abdomen. This
13 enables you to drag the casualty as you crawl.
14



15
16 *Figure B-10. Pistol-belt drag.*
17

18
19 (9) The *neck drag* (Figure B-11) is useful in combat because the bearer can
20 transport the casualty as he creeps behind a low wall or shrubbery, under a vehicle, or through a
21 culvert. If the casualty is unconscious, his head must be protected from the ground. The neck
22 drag cannot be used if the casualty has a broken arm.
23

24 **NOTE**

25 If the casualty is conscious, he may clasp his hands
26 together around your neck.
27

28 (a) Tie the casualty's hands together at the wrists.

29
30 (b) Straddle the casualty in a kneeling face-to-face position.

31
32 (c) Loop the casualty's tied hands over and around your neck.

33
34 (d) Crawl forward dragging the casualty with you.

1
2
3
4

NOTE

If the casualty is unconscious, protect his head from the ground.



5
6
7
8

Figure B-11. Neck drag.

9
10
11

(10) The *cradle-drop drag* (Figure B-12) is effective in moving a casualty up or down steps.

12
13
14

(a) Kneel at the casualty's head (with him lying on his back). Slide your hands, with palms up, under the casualty's shoulders and get a firm hold under his armpits.

15
16
17

(b) Rise (partially), supporting the casualty's head on one of your forearms. You may bring your elbows together and let the casualty's head rest on both of your forearms.

18
19
20

(c) Rise and drag the casualty backward. The casualty is in a semisitting position.

21
22
23

(d) Back down the steps, supporting the casualty's head and body and letting his hips and legs drop from step to step.

24
25

NOTE


If the casualty needs to be moved up the steps, you should back up the steps, using the same procedure.

26
27
28



1
2
3

Figure B-12. Cradle-drop drag.



4
5
6
7

Figure B-12. Cradle-drop drag (continued).

1 (11) The *LBE carry using the bearer's LBE* can be used with a conscious
2 casualty (Figure B-13).

3
4 (a) Loosen all suspenders on your LBE.


5
6 (b) Have the casualty place one leg into the loop formed by your
7 suspenders and pistol belt.

8
9 (c) Squat in front of the standing casualty. Have him place his other
10 leg into the loop, also.


11
12 (d) Have the casualty place his arms over your shoulders, lean forward
13 onto your back, and lock his hands together.

14
15 (e) Stand up and lean forward into a comfortable position.

16
17 (f) Continue the mission.
18



19
20 *Figure B-13. Load bearing equipment carry using bearer's LBE*
21 *(conscious casualty).*
22



1
2
3
4

Figure B-13. Load bearing equipment carry using bearer's LBE (conscious casualty) (continued).



Figure B-13. Load bearing equipment carry using bearer's LBE
(conscious casualty) (continued).

1
2
3
4
5 (12) The LBE carry using the bearer's LBE can be used with an unconscious
6 casualty or one who cannot stand (Figure B-14).

- 7
8 (a) Position the casualty on the flat of his back.
9
10 (b) Remove your LBE and loosen all suspender straps.
11
12 (c) Lift the casualty's leg and place it through the loop formed by your
13 suspenders and pistol belt. Then place the other leg. The LBE is moved up until the pistol belt is
14 behind the casualty's thighs.
15
16 (d) Lay between the casualty's legs; work his arms through his LBE
17 suspenders.
18
19 (e) Grasp the casualty's hand (on the injured side), and roll the
20 casualty (on his uninjured side) onto his back.

1
2
3
4
5
6
7
8
9

(f) Rise to one knee and then push into a standing position.


(g) Bring the casualty's arms over your shoulders. Grasp his hands and secure them if the casualty is unconscious. If the casualty is conscious, have him lock his hands in front if he is able to do so.

(h) Lean forward into a comfortable position and continue the mission.



10
11
12
13


Figure B-14. Load bearing equipment carry using bearer's LBE (unconscious casualty or one that cannot stand).



1
2 *Figure B-14. Load bearing equipment carry using bearer's LBE (unconscious*
3 *casualty or one that cannot stand [continued]).*
4
5

6 (13) The *LBE carry using the casualty's LBE* (Figure B-15) can be used with a
7 conscious or unconscious casualty.
8

- 1 (a) Position the casualty on his back with his LBE on.
2
3 (b) Loosen the casualty's two front suspenders.
4
5 (c) Position yourself between the casualty's legs, and slip your arms
6 into the casualty's two front suspenders (up to his shoulders).
7
8 (d) Work his arms out of his LBE suspenders.
9
10 (e) Grasp the casualty's hand (on the injured side), and roll him (on his
11 uninjured side) onto his back.
12
13 (f) Rise to one knee, then into a standing position.
14
15 (g) If the casualty is unconscious, grasp his hands and secure them,. If
16 the casualty is conscious, have him lock his hands in front of you.
17
18 (h) Lean forward into a comfortable position and continue the mission.
19



20
21 *Figure B-15. Load-bearing equipment carry using casualty's LBE.*
22



Figure B-15. Load-bearing equipment carry using casualty's LBE (continued).

1
2
3
4
5 **b. Two-Man Carries.** These carries should be used whenever possible. They
6 provide more casualty comfort, are less likely to aggravate injuries, and are less tiring for the
7 bearers. Five different two-man carries can be used.

8
9 (1) The *two-man support carry* (Figure B-16) can be used in transporting
10 either conscious or unconscious casualties. If the casualty is taller than the bearers, it may be
11 necessary for the bearers to lift the casualty's legs and let them rest on their forearms. The
12 bearers—
13

1
2 around his waist.

(a) Help the casualty to his feet and support him with their arms

3
4
5

(b) Grasp the casualty's wrists and draw his arms around their necks.



6
7
8
9

Figure B-16. Two-man supporting carry.

10 (2) The *two-man arms carry* (Figure B-17) is useful in carrying a casualty for
11 a moderate distance (50 to 300 meters) and placing him on a litter. To lessen fatigue, the bearers
12 should carry the casualty high and as close to their chests as possible. In extreme emergencies
13 when there is no time to obtain a spine board, this carry is the safest one for transporting a
14 casualty with a back injury. If possible, two additional bearers should be used to keep the
15 casualty's head and legs in alignment with his body. The bearers—

16

17 (a) Kneel at one side of the casualty and extend his arms above his
18 head; then they place their arms beneath the casualty's back, waist, hips, and knees.

19

20 (b) Lift the casualty while rising to their knees.

21

22 (c) Turn the casualty toward their chests, while rising to a standing
23 position. Carry the casualty high to lessen fatigue.

24




Figure B-17. Two-man arms carry.

1
2
3
4
5
6 (3) The *two-man fore-and aft-carry* (Figure B-18) is a useful *two-man carry*
7 for transporting a casualty for a long distance (over 300 meters). The taller of the two bearers
8 should position himself at the casualty's head. By altering this carry so that both bearers face the
9 casualty, it is useful for placing a casualty on a litter.

10
11 (a) One bearer spreads the casualty's legs and kneels between them
12 with his back to the casualty. He positions his hands behind the casualty's knees. The other
13 bearer kneels at the casualty's head, slides his hands under the arms, across the chest, and locks
14 his hands together.

15
16 (b) The two bearers rise together, lifting the casualty.
17



Figure B-18. Two-man fore-and-aft carry.

(4) Only a conscious casualty can be transported with the *four-hand seat carry* (Figure B-19) because he must help support himself by placing his arms around the bearers' shoulders. This carry is especially useful in transporting a casualty with a head or foot injury for a moderate distance (50 to 300 meters). It is also useful for placing a casualty on a litter.

(a) Each bearer grasps one of his wrists and one of the other bearer's wrists, thus forming a packsaddle.

(b) The two bearers lower themselves sufficiently for the casualty to sit on the packsaddle; then, they have the casualty place his arms around their shoulders for support. The bearers then rise to an upright position.




Figure B-19. Four-hand seat carry.

1
2
3
4
5
6
7

(5) The *two-hand seat carry* (Figure B-20) is used when carrying a casualty to a litter. With the casualty lying on his back, a bearer kneels on each side of the casualty at his hips. Each bearer passes his arms under the casualty's thighs and back, and grasps the other bearer's wrists. The bearers rise lifting the casualty.




Figure B-20. *Two-hand seat carry.*

8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28

B-9. Improvised Litters

Two men can support or carry a casualty without equipment for only short distances. By using available materials to improvise equipment, the casualty can be transported greater distances by two or more rescuers.

a. There are times when a casualty may have to be moved and a standard litter is not available. The distance may be too great for manual carries or the casualty may have an injury (such as a fractured neck, back, hip, or thigh) that would be aggravated by manual transportation. In these situations, litters can be improvised from materials at hand. Improvised litters must be as well constructed as possible to avoid risk of dropping or further injuring the casualty. Improvised litters are emergency measures and must be replaced by standard litters at the first opportunity.

b. Many different types of litters can be improvised, depending upon the materials available. A satisfactory litter can be made by securing poles inside such items as a blanket, poncho, shelter half, tarpaulin, mattress cover, jacket, shirt, or bed ticks, bags, and sacks. Poles


1 can be improvised from strong branches, tent supports, skis, lengths of pipe, or other objects. If
2 objects for improvising poles are not available, a blanket, poncho, or similar item can be rolled
3 from both sides toward the center so the rolls can be gripped for carrying a patient. Most flat-
4 surface objects of suitable size can be used as litters. Such objects include doors, boards,
5 window shutters, benches, ladders, cots, and chairs. If possible, these objects should be padded
6 for the casualty's comfort.

7
8 (1) To improvise a litter using a blanket and poles (Figure B-21), the
9 following steps should be used.

10
11 (a) Open the blanket and lay one pole lengthwise across the center;
12 then fold the blanket over the pole.

13
14 (b) Place the second pole across the center of the folded blanket.

15
16 (c) Fold the free edges of the blanket over the second pole and across
17 the first pole.
18



19
20 *Figure B-21. Litter made with blanket and poles.*
21
22

23 (2) To improvise a litter using shirts or jackets (Figure B-22), button the shirt
24 or jacket and turn it inside out, leaving the sleeves inside, (more than one shirt or jacket may be
25 required), then pass the pole through the sleeves.
26




Figure B-22. Litter improvised from jackets and poles.

(3) To improvise a litter from bed sacks and poles (Figure B-23), rip open the corners of bed ticks, bags, or sacks; then pass the poles through them.




Figure B-23. Litter improvised from bed sacks and poles.

(4) If no poles are available, roll a blanket, shelter half, tarpaulin, or similar item from both sides toward the center (Figure B-24). Grip the rolls to carry the casualty.




Figure B-24. Rolled blanket used as a litter.

1
2
3
4
5
6 *f.* Any of the appropriate carries may be used to place a casualty on a litter. These
7 carries are—

- 8
9
10
11
12
13
14
15
16
17
18
- The arms carry (Figure B-6).
 - The two-man arms carry (Figure B-17).
 - The two-man fore-and-aft carry (Figure B-18).
 - The two-hand seat carry (Figure B-2015).
 - The four-hand seat carry (Figure B-19).

19
20

WARNING

21 Unless there is an immediate life-threatening situation (such as
22 fire, explosion), **DO NOT** move the casualty with a suspected back
23 or neck injury. Seek medical personnel for guidance on how to
24 transport.

25 *g.* Either two or four persons (head/foot) may be used to lift a litter. To lift the litter,
26 follow the procedure below.

- 27
28
29
30
- (1) All carriers/bearers raise the litter at the same time.
 - (2) Keep the casualty as level as possible.

31
32

NOTE


33 Use caution when transporting on a sloping incline/hill.
34

APPENDIX C

DIGITAL PRESSURE

1
2
3
4
5
6
7
8
9
10
11
12
13
14

Digital pressure (often called "pressure points") is an alternate method to control bleeding. This method uses pressure from the fingers, thumbs, or hands to press at the site or point where a main artery supplying the wounded area lies near the skin surface or over bone (Figure C-1). This pressure may help shut off or slow down the flow of blood from the heart to the wound and is used in combination with direct pressure and elevation. It may help in instances where bleeding is not easily controlled, where a pressure dressing has not yet been applied, or where pressure dressings are not readily available.



15
16
17

Figure C-1. Digital pressure (pressure with fingers, thumbs or hands).

GLOSSARY

ABBREVIATIONS AND ACRONYMS

AC	hydrogen cyanide
AR	Army regulation
ATM	advanced trauma management
BDU	battle dress uniform
BZ	anticholinergic drugs
C	Celsius
cc	cubic centimeter
CANA	Convulsant Antidote for Nerve Agent
CG	phosgene
CHS	combat health support
CK	cyanogen chloride
Cl	chlorine
CNS	central nervous system
CO ₂	carbon dioxide
CSR	combat stress reaction
CTA	common table of allowance
CX	phosgene oxime
DA	Department of the Army
DM	(adamsite) diphenylaminochloroarsine
DNBI	disease and nonbattle injury
DOD	Department of Defense
DP	diphosgene
DS	direct support
EMT	emergency medical treatment
F	Fahrenheit
FM	field manual
H	mustard
HD	mustard
HN	nitrogen mustard
IV	intravenous
JAMA	Journal of the American Medical Association
L	lewisite
LBE	load bearing equipment
MILES	Multiple Integrated Laser Engagement System
ml	milliliter
MOPP	mission-oriented protective posture
MOS	military occupational specialty
MTF	medical treatment facility
NAPP	Nerve Agent Pyridostigmine Pretreatment
NATO	North Atlantic Treaty Organization

FM 21-11
Initial Draft

NBC	nuclear, biological, and chemical
oz	ounce
2 PAM Cl	pralidoxime chloride
PAM	pamphlet
PS	chloropicrin
PTSD	post-traumatic stress disorder
STANAG	standardization agreement
STP	soldier training publication
TB MED	technical bulletin medical
TM	technical manual
TSOP	tactical standing operating procedure
US	United States
WP	white phosphorus

REFERENCES

DOCUMENTS NEEDED

These documents must be available to the intended users of this publication.

Joint and Multiservice Publications

FM 21-10 Field Hygiene and Sanitation (AFM 161-10)

Army Publications

AR 600-110 Identification, Surveillance and Administration of Personnel Infected with Human Immunodeficiency Virus (HIV)

DA PAM 351-20 Army Correspondence Course Program Catalog

FM 3-4 NBC Protection

FM 3-5 NBC Decontamination

FM 3-100 NBC Operations

FM 8-10-6 Medical Evacuation in a Theater of Operations—Tactics, Techniques, and Procedures

FM 8-10-7 Health Service Support in a Nuclear, Biological, and Chemical Environment

FM 8-35 Evacuation of the Sick and Wounded

FM 8-230 Medical Specialist

FM 8-285 Treatment of Chemical Agent Casualties and Conventional Chemical Injuries

FM 22-51 Leaders' Manual for Combat Stress Control

FM 26-2 Management of Stress in Army Operations

CTA 8-100 Army Medical Department Expendable/Durable Items

CTA 50-900 Clothing and Individual Equipment

STP 21-1-SMCT Soldier's Manual of Common Tasks (Skill Level I)

READINGS RECOMMENDED

These readings contain relevant supplemental information.

Joint and Multiservice Publications

FM 8-9 NATO Handbook on the Medical Aspects of NBC Defensive Operations (AMED P-6; NAVMED P-5059; AFP 161-3)

FM 8-33 Control of Communicable Diseases in Man, 14th Edition (NAVMED P-5038)

TB MED 81 Cold Injury (NAVMED P-5052-29; AFP 161-11)

TB MED 507 Occupational and Environmental Health Prevention, Treatment and Control of Heat Injury (NAVMED P-5052-5; AFP 161-1)

TM 8-285 Treatment of Chemical Agent Casualties and Conventional Military Chemical Injuries (NAVMED P-5041; AFM 160-12)

FM 21-11
Final Draft

Army Publications

AR 310-25	Dictionary of United States Army Terms (Short Title AD) (Reprinted with basic including Change 1)
AR 310-50	Authorized Abbreviations, Brevity Codes, and Acronyms
DA PAM 40-12	Who Needs It--Venereal Diseases
TM 3-4230-216-10	Operator's Manual for Decontaminating Kit, Skin: M258A1 (NSN 4230-01-101-3984) and Training Aid, Skin Decontaminating: M58A1 (6910-01-101-1768)

Nonmilitary Publications:

American Heart Association. A Manual for Instructors of Basic Cardiac Life Support. Dallas: American Heart Association, 1981.

———. Instructors Manual for Basic Life Support. Dallas: American Heart Association, 1985.

JAMA. "Standards and Guidelines for Cardiopulmonary Resuscitation (CPR) and Emergency Cardiac Care (ECC)." The Journal of the American Medical Association, Vol 255, No. 21 (June 1986): 2905-2989.